

RESEARCH ARTICLE

Open Access

# Little impact of tsunami-stricken nuclear accident on awareness of radiation dose of cardiac computed tomography: A questionnaire study

Sung Hea Kim<sup>1</sup>, Hyun-Joong Kim<sup>1</sup>, Hyun Kyun Ki<sup>1</sup>, Eui-Jong Chung<sup>1</sup>, Soon Yong Suh<sup>2</sup>, Seong Woo Han<sup>3</sup> and Kyu-Hyung Ryu<sup>4\*</sup>

## Abstract

**Background:** With the increased use of cardiac computed tomography (CT), radiation dose remains a major issue, although physicians are trying to reduce the substantial risks associated with use of this diagnostic tool. This study was performed to investigate recognition of the level of radiation exposure from cardiac CT and the differences in the level of awareness of radiation before and after the Fukushima nuclear plant accident.

**Methods:** We asked 30 physicians who were undergoing training in internal medicine to determine the equivalent doses of radiation for common radiological examinations when a normal chest X-ray is accepted as one unit; questions about the absolute radiation dose of cardiac CT data were also asked.

**Results:** According to the results, 86.6% of respondents believed the exposure to be 1 mSv at most, and 93.3% thought that the exposure was less than that of 100 chest X-rays. This finding indicates that their perceptions were far lower than the actual amounts. Even after the occurrence of such a large nuclear disaster in Fukushima, there were no significant differences in the same subjects' overall awareness of radiation amounts.

**Conclusions:** Even after such a major social issue as the Fukushima nuclear accident, the level of awareness of the accurate radiation amount used in 64-channel multidetector CT (MDCT) by clinical physicians who order this test was not satisfactory. Thus, there is a need for the development of effective continuing education programs to improve awareness of radiation from ionizing radiation devices, including cardiac CT, and emphasis on risk-benefit evaluation based on accurate knowledge during medical training.

**Keywords:** 64-channel MDCT, Radiation dose, Questionnaires, Awareness

## Background

With recent advances in technology, the importance of cardiac imaging has also increased. In particular, the roles of both cardiac computed tomography (CT) and magnetic resonance imaging (MRI) have expanded. Cardiac CT, because of its advantage of allowing for noninvasive evaluation of both coronary artery stenosis and myocardial perfusion, has been suggested as a substitute tool for conventional coronary angiography in the early detection of ischemic cardiac disease [1,2]. Although

recent reports have recognized its limitation in replacement of conventional coronary angiography in terms of diagnostic accuracy of coronary artery diseases, the frequency of use of cardiac CT in clinical practice is gradually increasing [3,4].

However, unlike MRI, cardiac CT has an innate issue with regard to its radiation hazard, which is known to be higher than that of other CT modalities because of cardiac motion. For proper use of cardiac CT, clinicians must have accurate knowledge of not only its advantages, but also its limitations, especially with regard to radiation hazard. Thus, awareness of radiation hazard by healthcare professionals is essential and important for proper evaluation of the risk-benefit ratio.

\* Correspondence: khryumd@hanmail.net

<sup>4</sup>Division of Cardiology, Department of Internal Medicine, College of Medicine, Hallym University Dongtan Sacred Heart Hospital, 40, Sukwoo-Dong, Hwaseong-Si, Gyeonggi-Do 445-170, South Korea  
Full list of author information is available at the end of the article


The aim of this study was to investigate medical residents' awareness of the radiation dose used in the performance of cardiac CT and assess the impact of social issues, such as the Japan nuclear plant explosion, on the level of awareness.

### Methods

The IRB exempted survey was conducted from November 2009 to March 2010 as part of the training and assessment of medical residents (KUH 1010479). The survey aimed to improve their perceptions of the absolute radiation dose in one prescription of 64-channel multidetector (MDCT) and the relative radiation dose compared with that of a simple chest PA. The subjects included 30 medical residents who were undergoing training in internal medicine. An earthquake subsequently occurred in Fukushima, Japan in March 2011. A follow-up survey was performed on the same subjects using the same questionnaire. The change in the degree of awareness of pre- and post-earthquake radiation exposure was evaluated. Research carried out in compliance with the Helsinki Declaration.

The survey sheet included two categories: the general characteristics of the respondents and the awareness of the exposure by each test. Regarding awareness of radiation exposure, the first survey asked questions on radiation doses for conventional angiography, cardiac MDCT, and MIBI scan compared with a simple chest PA to determine the relative radiation dose. In the second questionnaire, subjects were asked to determine the absolute radiation dose for cardiac MDCT (Additional file 1: Figure S1).

Numerical variables are shown as mean  $\pm$  standard deviation or median (minimum, maximum) values, and categorical variables are shown as number and percentage.



The Kolmogorov-Smirnov test was used to determine normal distribution. Nonparametric statistics, including the Wilcoxon rank sum test, were employed for comparison of differences in variables between the first and second survey;  $p < 0.05$  was considered statistically significant. PASW statistics software (version 17) was used to perform the statistical analysis.

### Results


The mean age of the subjects was  $31 \pm 2.5$  years, and 19 subjects (63.3%) were male. The median value of the absolute radiation exposure was 1 mSv (0.5 mSv, 100 mSv) on the first survey and 1 mSv (0.1 mSv, 300 mSv) on the second survey. According to the results of the Wilcoxon rank sum test, there was no difference between the first and second survey ( $z = -1.483$ ,  $p = 0.138$ ).

On the first survey, the median value of the relative radiation exposure of cardiac CT compared with a simple chest X-ray was 50 times (10, 500), which did not differ from that of the second survey (50 times [10, 1000]) ( $z = -4.47$ ,  $p = 0.655$ ).

The 64-channel MDCT is generally known to carry a radiation exposure of 5 to 10 mSv per shot, which is equivalent to 300 times the amount of a simple chest X-ray. However, 86.6% of the respondents on the first survey and 83.3% on the second survey perceived the exposure to be 1 mSv at most. On the first and second surveys, 93.3% and 90.0% of respondents thought that the exposure was 100 times less than the amount in the chest X-ray, respectively (Figures 1 and 2).

### Discussion

In our study, we found that medical residents' level of knowledge of the radiation dose of cardiac CT is low and inadequate. However, previous studies have highlighted the lack of appreciation of radiation doses,



even in radiologists [5-7]; therefore, this result is not surprising.

Second, there was no improvement in the clinicians' degree of awareness regarding radiation exposure between the pre- and post-earthquake periods, despite the experience of such a major social issue and the resulting information campaign about exposure through newspapers and television for several weeks, which caused even laymen to become familiar with the international unit of radiation dose (the sievert). This may suggest that the clinicians believed that their underestimated radiation dose was adequate and not in need of correction, even after the nuclear plant accident, which indicates a fixed underestimated status.

As the clinical areas of CT expand, the frequency of use of CT and concerns about increases in the exposure dose are likewise increasing. However, even a small exposure dose could be hazardous, as seen during the Hiroshima atomic bombing [8,9]; therefore, clinicians should make every effort to minimize the radiation dose.

To decrease the radiation dose from CT equipment used in cardiac MDCT, efforts have been made to control the amount of X-ray by the attenuation level or to decrease the amount of X-ray at the systolic phase of electrocardiography (ECG); these efforts have resulted in some verifiable success. One study reported on a trial to decrease the radiation exposure by up to 5 mSv using a low-dose coronary CT technique with 100 kVp rather than 120 kVp; however, the diagnostic accuracy of this technique has not yet been verified [10]. Various strategies to decrease the radiation dose in coronary CT angiography or coronary calcium CT have been applied, including the use of low amperage, low voltage, and ECG-triggered amperage modulation. In particular, ECG-triggered amperage modulation has been reported to lower the effective dose by 20% to 50% based on the heart rate [10-13]. However, prior to the establishment of technical advancements, improvements in clinicians' awareness and knowledge of radiation should occur, because proper knowledge regarding risk is the first step in risk-benefit analysis in our practice.

There could be several explanations for the low level of clinicians' awareness of radiation. Clinicians' focus on treatment methods and the relative indifference to the test method itself may play a major role in the lack of understanding. Other reasons for the lack of understanding may include the image-centric thinking pattern of the modern generation, lack of systematic and repetitive education systems, and lack of knowledge of radiation hazard.

As demonstrated in this study, a one-time shock has no significant educational effect. Thus, systematic and repetitive education systems may be first to be established ultimately to increase awareness of the potential risk of

radiation. Another way to increase awareness is visualization of total radiation doses using figures or graphics on its readings, not only on images, which may alarm clinicians to recall the risk and benefit once more [14].

### Study limitations

First, the number of subjects was small to represent all of trainee or medical residents. Second, our survey was not originally designed to compare the level of perceptions of radiation dose before and after social issue, rather to improve their perceptions of the radiation dose as part of the annual training program. Thus, the questionnaires used in our survey may have not sufficient validity and reliability to evaluate the awareness of radiation dose. But, because there is no established tool of evaluating the level of perceptions of radiation dose, the recent studies have to depend on questionnaire survey [15-20].

### Conclusions

The results of our study highlight that the level of awareness of the accurate radiation amount used in the performance of 64-channel MDCT by clinical physicians who order the test is not satisfactory. Therefore, it is suggested that clinical physicians can safely perform multiple-channel MDCT on patients not only through being properly informed on the use of multiple-channel MDCT, but also through active training on the possible risk of patients' exposure. In addition, development of an effective educative method and the optimum method for improvement of awareness is needed.

### Additional file

**Additional file 1: Figure S1.** Key questionnaires used in the Survey

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

SHK and EJC participated in data analysis and drafted the initial manuscript. HJK and SYS participated in the sequence alignment and critical revision. HKK and SWH participated in the design of the study and performed the statistical analysis. KHR conceived the study, participated in its design and coordination, and helped to draft the manuscript. All authors made substantial contributions to the acquisition of data. All authors read and approved the final manuscript.

### Author details

<sup>1</sup>Department of Internal Medicine, Konkuk University School of Medicine, 1, Hwayang-dong, Gwangjin-gu, Seoul, South Korea. <sup>2</sup>Heart Center, Gachon University of Medicine and Science, Gil Hospital, Incheon, South Korea. <sup>3</sup>Department of Internal Medicine, College of Medicine, Hallym University, Seoul, South Korea. <sup>4</sup>Division of Cardiology, Department of Internal Medicine, College of Medicine, Hallym University Dongtan Sacred Heart Hospital, 40, Sukwoo-Dong, Hwaseong-Si, Gyeonggi-Do 445-170, South Korea.

Received: 7 December 2011 Accepted: 18 April 2013  
Published: 30 April 2013

## References

1. Yoo SM, Rho JY, Lee HY, Song IS, Moon JY, White CS: **Current concepts in cardiac CT angiography for patients with acute chest pain.** *Korean Circ J* 2010, **40**:543–549.
2. Gaudio C, Mirabelli F, Pelliccia F, Francone M, Tanzilli G, Di Michele S, Leonetti S, De Vincentis G, Carbone I, Mangieri E, Catalano C, Passariello R: **Early detection of coronary artery disease by 64-slice multidetector computed tomography in asymptomatic hypertensive high-risk patients.** *Int J Cardiol* 2009, **135**:280–286.
3. Miller JM, Rochitte CE, Dewey M, Arbab-Zadeh A, Niinuma H, Gottlieb L, Paul N, Clouse ME, Shapiro EP, Hoe J, Lardo AC, Bush DE, de Roos A, Cox C, Brinker J, Lima JA: **Diagnostic performance of coronary angiography by 64-row CT.** *N Engl J Med* 2008, **359**:2324–2336.
4. Paech DC, Weston AR: **A systematic review of the clinical effectiveness of 64-slice or higher computed tomography angiography as an alternative to invasive coronary angiography in the investigation of suspected coronary artery disease.** *BMC Cardiovasc Disord* 2011, **11**:32.
5. Lee RK, Chu WC, Graham CA, Rainer TH, Ahuja AT: **Knowledge of radiation exposure in common radiological investigations: a comparison between radiologists and non-radiologists.** *Emerg Med J* 2012, **29**:306–308.
6. Zhou GZ, Wong DD, Nguyen LK, Mendelson RM: **Student and intern awareness of ionising radiation exposure from common diagnostic imaging procedures.** *J Med Imaging Radiat Oncol* 2010, **54**:17–23.
7. Soye JA, Paterson A: **A survey of awareness of radiation dose among health professionals in Northern Ireland.** *Br J Radiol* 2008, **81**:725–729.
8. Pierce DA, Preston DL: **Radiation-related cancer risks at low doses among atomic bomb survivors.** *Radiat Res* 2000, **154**:178–186.
9. Preston DL, Ron E, Tokuoka S, Funamoto S, Nishi N, Soda M, Mabuchi K, Kodama K: **Solid cancer incidence in atomic bomb survivors: 1958–1998.** *Radiat Res* 2007, **168**:1–64.
10. Abada HT, Larchez C, Daoud B, Sigal-Cinqualbre A, Paul JF: **MDCT of the coronary arteries: feasibility of low-dose CT with ECG-pulsed tube current modulation to reduce radiation dose.** *Am J Roentgenol* 2006, **186**(6 Suppl 2):387–390.
11. Shemesh J, Evron R, Koren-Morag N, Apter S, Rozenman J, Shaham D, Itzchak Y, Motro M: **Coronary artery calcium measurement with multidetector row CT and low radiation dose: comparison between 55 and 165 mAs.** *Radiology* 2005, **236**:810–814.
12. Jakobs TF, Becker CR, Ohnesorge B, Flohr T, Suess C, Schoepf UJ, Reiser MF: **Multi-slice helical CT of the heart with retrospective ECG gating: reduction of radiation exposure by ECG-controlled tube current modulation.** *Eur Radiol* 2002, **12**:1081–1086.
13. Jung B, Mahnken AH, Stargardt A, Simon J, Flohr TG, Schaller S, Koos R, Gunther RW, Wildberger JE: **Individually weight-adapted examination protocol in retrospectively ECG-gated MSCT of the heart.** *Eur Radiol* 2003, **13**:2560–2566.
14. Sung DW, et al: **Radiation exposure in diagnostic areas: issues and countermeasures.** *J Korean Med Assoc* 2011, **54**:1246–1247.
15. Merzenich H, Krille L, Hammer G, Kaiser M, Yamashita S, Zeeb H: **Paediatric CT scan usage and referrals of children to computed tomography in Germany—a cross-sectional survey of medical practice and awareness of radiation related health risks among physicians.** *BMC Health Serv Res* 2012, **12**:47.
16. Puri S, Hu R, Quazi RR, Voci S, Veazie P, Block R: **Physicians' and midlevel providers' awareness of lifetime radiation-attributable cancer risk associated with commonly performed CT studies.** *Am J Roentgenol* 2012, **199**:1328–1336.
17. Brown N, Jung JL: **Knowledge of medical imaging radiation dose and risk among doctors.** *J Med Imaging Radiat Oncol* 2013, **57**(1):8–14.
18. Uri IF, et al: **Lack of radiation awareness among referrers: implications and possible solutions.** *Int J Clin Pract* 2012, **66**(6):574–581.
19. Divrik Gökçe S, Gökçe E, Coşkun M: **Radiology residents' awareness about ionizing radiation doses in imaging studies and their cancer risk during radiological examinations.** *Koreab J Radiol* 2012, **13**(2):202–209.
20. Sin HK, Wong CS, Huang B, Yiu KL, Wong WL, Chu YC: **Assessing local patients' knowledge and awareness of radiation dose and risks associated with medical imaging.** *J Med Imaging Radiat Oncol* 2013, **57**(1):38–44.

doi:10.1186/1756-0500-6-170

**Cite this article as:** Kim et al.: Little impact of tsunami-stricken nuclear accident on awareness of radiation dose of cardiac computed tomography: A questionnaire study. *BMC Research Notes* 2013 **6**:170.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

