


DATA NOTE

Open Access



Draft genome sequences of *Mycolicibacterium peregrinum* isolated from a pig with lymphadenitis and from soil on the same Japanese pig farm

Tetsuya Komatsu¹, Kenji Ohya^{2,3,4,10}, Kotaro Sawai^{2,5}, Justice Opare Odoi³, Keiko Otsu⁶, Atsushi Ota⁷, Toshihiro Ito^{8,11}, Mikihiro Kawai⁸ and Fumito Maruyama^{8,9*} 

Abstract

Objectives: *Mycolicibacterium peregrinum*, a rapidly growing mycobacterial species, can opportunistically infect humans and other animals. Although *M. peregrinum* infections in animals have been reported, the infection sources are unknown, as is information on its virulence and drug resistant genes, which limits our current understanding of this bacterium. To address this knowledge gap, we obtained draft genome sequences for two *M. peregrinum* isolates; one from a case of pig lymphadenitis and one from the pig farm's soil.

Data description: We report here the draft genome sequences of *M. peregrinum* isolates 131_1 and 138 (6,451,733-bp and 6,479,047-bp). They were isolated from a pig with mesenteric lymph node lymphadenitis and from soil on the Japanese farm where the pig was reared. A sequence alignment identity score of 100% was obtained by in silico DNA–DNA hybridization of the two isolates, while 98.28% (isolate 131_1) and 98.27% (isolate 138) scores were recorded for hybridization with a human isolate. Both isolates carry *arr-1*, *AAC(2′)-Ib*, *RbpA*, *mtrA* and *tap* drug-resistance genes. Isolates 131_1 and 138 carry 234 and 236 putative virulence genes, respectively. Therefore, environment *M. peregrinum* is potentially drug resistant and can cause swine lymphadenitis. Our data provides valuable new information for future studies on nontuberculous mycobacteria.

Keywords: *Mycolicibacterium peregrinum*, Draft genome sequence, Mycobacterium, Drug resistance, Virulence, In silico DNA–DNA hybridization, Pig, Pig farm

Objective

Mycolicibacterium peregrinum (basonym: *Mycobacterium peregrinum*), a known pathogenic and rapidly growing mycobacterium (RGM), has been isolated from clinical samples from pigs, cattle and a person [1–3]. Several cases of *M. peregrinum* infection have been reported in aquatic animals [4, 5], wild animals [6–8] and livestock [1, 2, 9], including one porcine case [1]. Nontuberculous mycobacteria (NTM) such as *M. peregrinum* generally

reside in water and soil, and these environmental NTM are believed to occasionally infect humans and other species, opportunistically [10]. However, the transmission sources for *M. peregrinum* in humans and other animals are not clear in each case. Classification of the *Mycobacteria* genus currently positions the *Mycobacterium fortuitum* group, including *M. peregrinum*, as *Mycolicibacterium* [11].

Few studies on *M. peregrinum* virulence genes have been conducted [12], but the medical fields have reported on multidrug resistance in this bacterium [13]. It has also been reported that *M. peregrinum* is more susceptible to some antimicrobial agents than other mycobacteria species [14]. Other studies have reported that some RGM

*Correspondence: maruyama.fumito.5e@kyoto-u.ac.jp

⁸ Department of Microbiology, Graduate School of Medicine, Kyoto University, Yoshida-Konoe-cho, Sakyo-ku, Kyoto 6068501, Japan
Full list of author information is available at the end of the article



carry antibiotic resistance genes, such as erythromycin ribosomal methylase (*erm*) [15], *LfrA* and *tap* [16]. Although the *tap* gene is present in *M. peregrinum*, a comprehensive analysis of its antibiotic resistance genes has not been done. Therefore, to obtain better understanding of the potential risk posed by antibiotic resistance in *M. peregrinum*, an analysis at the draft genome level is necessary. Such information would be useful to veterinary medicine as there is no genome information on isolates from non-human animals. To aid future investigations into the sources of *M. peregrinum* infection and to provide information on virulence and drug resistance genes, we present here the draft *M. peregrinum* genome sequences for isolates 131_1 and 138 from a case of swine lymphadenitis and from soil on the same Japanese farm, respectively.

Data description

Mycobacterium peregrinum isolate 131_1 was isolated from the mesenteric lymph nodes of a pig with lymphadenitis and isolate 138 was isolated from soil on the same pig farm (Tokai area of Japan), as described previously [17]. Both samples were individually decontaminated with an equal volume of 2% NaOH and then inoculated onto 2% Ogawa medium (Kyokuto Pharmaceutical, Tokyo, Japan). Both isolates were species identified by sequencing the 16S rRNA, *hsp65* and *rpoB* genes [18, 19]. Genomic DNA was extracted using the Pure-Link genomic DNA extraction kit (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions, and paired-end libraries with an average insert size of 350-bp were prepared. Sequencing (2 × 150-bp) was conducted on the HiSeq X Ten sequencing platform (Illumina, San Diego, CA, USA) at the Beijing Genomics Institute (Shenzhen, China). Draft genome sequences were obtained from the reads according to the method reported previously (Table 1) [17]. In brief, the reads were trimmed by TrimGalore! (<https://github.com/FelixKrueger/TrimGalore>) and mismatched reads were corrected, assembled and polished using SPAdes [20], Pilon [21] and Unicycler [22]. Genome completeness was estimated using CheckM [23]. Taxonomic classification was conducted using Kaiju [24] and Anvi'o [25]. Draft genomes were annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) [26]. Virulence

and drug resistant genes were identified by VFanalyzer (<http://www.mgc.ac.cn/VFs/main.htm>) and RGI (<https://card.mcmaster.ca/analyze/rgi>). In silico DNA–DNA hybridization was conducted by the MUMmer program with JspiecesWS [27].

The draft genome sequence of *M. peregrinum* isolate 131_1 (Data file 1) comprised 33 contigs with a total length of 6,451,733 bp, a G+C content of 66.41%, and an N50 size of 292,445 bp. The *M. peregrinum* 138 isolate's draft genome sequence (Data file 2) comprised 46 contigs with a total length of 6,479,047 bp, a G+C content of 66.41%, and an N50 size of 324,444 bp. The coding sequences, rRNAs and tRNAs in both isolates were estimated at 6169, 3, and 55 (isolate 131_1) and 6180, 3, and 55 (isolate 138), respectively. Both isolates contained large numbers of putative virulence genes and genes involved in metabolism (e.g., amino acid, purine, lipid and fatty acid genes), anaerobic respiration, anti-apoptosis, catabolism, metal uptake, cell surface components, mammalian cell entry operons, phagosome arrest, proteases, regulation, secreted proteins, secretion system, stress adaptation and toxins. Both isolates contain five drug resistance-related genes: *arr-1*, *AAC(2')-Ib*, *RbpA*, *mtrA* and *tap*. In silico DNA–DNA hybridization revealed that the aligned nucleotide sequences from *M. peregrinum* isolates 131_1 and 138 share 98.28% and 98.27% identity with the human *M. peregrinum* isolate [3], respectively, 88.46% sequence identity with *M. fortuitum* subsp. *fortuitum* [28], 85.18% sequence identity with *Mycobacteroides abscessus* [29], 84.60% and 84.61% identity with *M. mucogenicum* [30], respectively, 84.50% sequence identity with *Mycobacteroides chelonae* [31], and 84.21% sequence identity with *M. neoaurum* [32]. An aligned sequence identity score of both isolates was 100%, suggesting that *M. peregrinum* exists in the farm soil and both isolates might possibly be the same origin. Sequencing revealed that both isolates may be resistant to rifampin and macrolide antibiotics. These results provide useful information for future NTM studies and for clinical antibiotic use.

Limitations

The present data are based on the genome sequences of *M. peregrinum* isolates 131_1 and 138 at the draft level. Therefore, the exact lengths of these sequences, numbers

Table 1 Overview of data files

Label	Name of data file/data set	File types (file extension)	Data repository and identifier (DOI or accession number)
Data file 1	<i>M. peregrinum</i> 131_1	FASTA file (.fasta)	GenBank RWJZ00000000 (https://www.ncbi.nlm.nih.gov/nucore/RWJZ00000000)
Data file 2	<i>M. peregrinum</i> 138	FASTA file (.fasta)	GenBank RWKA00000000 (https://www.ncbi.nlm.nih.gov/nucore/RWKA00000000)

of coding sequences, rRNAs, tRNAs and repetitive elements cannot be predicted with certainty. The existence of plasmid/s or extra-chromosomal DNAs also cannot be predicted with certainty.

Abbreviations

LfrA: the membrane efflux pump gene for quinolones (confers resistance to macrolides); *erm*: ribosomal RNA methyltransferase gene; *tap*: major facilitator superfamily (MFS) antibiotic efflux pump gene (confers resistance to tetracyclines); *arr-1*: rifampin ADP-ribosyltransferase (Arr) gene; *AAC(2)-Ib*: chromosomal-encoded aminoglycoside acetyltransferase gene (confers resistance to aminoglycosides); *RbpA*: RNA-polymerase binding protein gene (confers resistance to rifampin); *mtrA*: transcriptional activator gene of the MtrCDE multidrug efflux pump (confers resistance to penam, a macrolide antibiotic).

Acknowledgements

We thank the Data Integration and Analysis Facility at the National Institute for Basic Biology for providing some of the computational resources. We also thank Sandra Cheesman, PhD, from Edanz Group (<http://www.edanzediting.com/ac>) for editing a draft of this manuscript.

Authors' contributions

KOh and FM designed the research and conceived the experiments. KOt collected the samples and took part in the bacterial isolation experiments. TK, KOh, KS, JOO, AO, TI, MK and FM conducted the experiments and analyzed the data. TK, KOh and FM wrote the manuscript. All authors read and approved the final manuscript.

Funding

This study was supported by a grant from the Japan Agency for Medical Research and Development (AMED) (17fk0108116h0401), the Japan Racing Association (JRA) Livestock Industry Promotion Project (H28-29_239, H29-30_7) of the JRA, a grant for Meat and Meat Products (H28-130, H30-60) managed by the Ito Foundation for Research in design study, collection, analysis; and was supported by grants from the Japan Society for the Promotion of Science (JSPS) KAKENHI (JP26304039, JP18K19674, 16H05501, 16H01782) and the Joint Research Program of the Research Center for Zoonosis Control in Hokkaido University in interpretation of data, writing the manuscript, and English editing. JOO is a recipient of a Japanese government (MEXT) scholarship.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Availability of data materials

The data described in this Data note can be freely and openly accessed on GenBank. Please see Table 1 and the reference list for details and links to the data accession: RWJZ00000000 (<https://www.ncbi.nlm.nih.gov/nuccore/RWJZ00000000>) and accession: RWKA00000000 (<https://www.ncbi.nlm.nih.gov/nuccore/RWKA00000000>).

Author details

¹ Aichi Prefectural Chuo Livestock Hygiene Service Center, 1-306 Jizouno, Miaicho, Okazaki 4440805, Japan. ² Laboratory of Veterinary Microbiology, Faculty of Applied Biological Sciences, Gifu University, 1-1 Yanagido, Gifu 5011193, Japan. ³ United Graduate School of Veterinary Sciences, Gifu University, 1-1 Yanagido, Gifu 5011193, Japan. ⁴ Education and Research Center for Food Animal Health, Gifu University (GeFAH), 1-1 Yanagido, Gifu 5011193, Japan. ⁵ Viral Disease and Epidemiology Research Division, National Institute of Animal Health, National Agriculture Research Organization, 3-1-5 Kannondai, Tsukuba, Ibaraki 3050856, Japan. ⁶ Gifu Prefectural Chuo Livestock Hygiene Service Center, 1-1 Yanagido, Gifu 5011112, Japan. ⁷ Data Science Center, Division of Biological Science, Nara Institute of Science and Technology, Ikoma, Nara

6300192, Japan. ⁸ Department of Microbiology, Graduate School of Medicine, Kyoto University, Yoshida-Konoe-cho, Sakyo-ku, Kyoto 6068501, Japan. ⁹ Scientific and Technological Bioresource Nucleus, Universidad de La Frontera, Av. Francisco Salazar 01145, Temuco 4811230, Chile. ¹⁰ Present Address: Division of Microbiology, National Institute of Health Sciences, 3-25-26 Tonomachi, Kawasaki-ku, Kawasaki 210-9501, Japan. ¹¹ Present Address: Laboratory of Proteome Research, Proteome Research Center, National Institute of Biomedical Innovation, Ibaraki, Osaka 567-0085, Japan.

Received: 3 May 2019 Accepted: 11 June 2019

Published online: 17 June 2019

References

- Cvetnić Z, Spicić S, Benić M, Katalinić-Janković V, Pate M, Krt B, Ocepek M. Mycobacterial infection of pigs in Croatia. *Acta Vet Hung.* 2007;55:1–9.
- Nuru A, Zewude A, Mohammed T, Wondale B, Teshome L, Getahun M, Mamo G, Medhin G, Pieper R, Ameni G. Nontuberculous mycobacteria are the major causes of tuberculosis like lesions in cattle slaughtered at Bahir Dar Abattoir, northwestern Ethiopia. *BMC Vet Res.* 2017;13:237.
- Omar SV, Allam M, Joseph L, Mtshali S, Ismail NA, Ismail A. Draft genome sequence of *Mycobacterium peregrinum* isolated from an HIV-positive patient in South Africa. *Genome Announc.* 2017;5:e00759–817.
- Mrlik V, Slany M, Kubecka J, Seda J, Necas A, Babak V, Slana I, Kriz P, Pavlik I. A low prevalence of mycobacteria in freshwater fish from water reservoirs, ponds and farms. *J Fish Dis.* 2012;35:497–504.
- Gcebe N, Michel AL, Hlokwé TM. Non-tuberculous *Mycobacterium* species causing mycobacteriosis in farmed aquatic animals of South Africa. *BMC Microbiol.* 2018;18:32.
- Vitali SD, Eden PA, Payne KL, Vaughan RJ. An outbreak of mycobacteriosis in Gouldian finches caused by *Mycobacterium peregrinum*. *Vet Clin North Am Exot Anim Pract.* 2006;9:519–22.
- de Lisle GW, Kawakami RP, Yates GF, Collins DM. Isolation of *Mycobacterium bovis* and other mycobacterial species from ferrets and stoats. *Vet Microbiol.* 2008;132:402–7.
- Liu H, Yan J, Luo J, Yan R, Chen H, Cheng H, Liu D, He H. Mycobacteriosis associated with *Mycobacterium peregrinum* infection in Red-crowned Cranes (*Grus japonensis*) in China. *J Wildl Dis.* 2014;50:703–6.
- Padya L, Chin'ombe N, Magwenzi M, Mbanga J, Ruhanya V, Nziramasanga P. Molecular identification of *Mycobacterium* species of public health importance in cattle in Zimbabwe by 16S rRNA gene sequencing. *Open Microbiol J.* 2015;31:38–42.
- Nishiuchi Y, Iwamoto T, Maruyama F. Infection sources of a common non-tuberculous mycobacterial pathogen, *Mycobacterium avium* complex. *Front Med (Lausanne).* 2017;4:27.
- Gupta RS, Lo B, Son J. Phylogenomics and comparative genomic studies robustly support division of the genus *Mycobacterium* into an emended genus *Mycobacterium* and four novel genera. *Front Microbiol.* 2018;9:67.
- Harriff MJ, Wu M, Kent ML, Bermudez LE. Species of environmental mycobacteria differ in their abilities to grow in human, mouse, and carp macrophages and with regard to the presence of mycobacterial virulence genes, as observed by DNA microarray hybridization. *Appl Environ Microbiol.* 2008;74:275–85.
- Santos A, Cremades R, Rodriguez JC, Garcia-Pachon E, Ruiz M, Royo G. *Mycobacterium peregrinum*: bactericidal activity of antibiotics alone and in combination. *J Infect Chemother.* 2008;14:262–3.
- Cavusoglu C, Gurpinar T, Ecemis T. Evaluation of antimicrobial susceptibilities of rapidly growing mycobacteria by Sensititre RAPMYCO panel. *New Microbiol.* 2012;35:73–6.
- Brown-Elliott BA, Hanson K, Vasiredy S, Iakhiaeva E, Nash KA, Vasiredy R, Parodi N, Smith T, Gee M, Strong A, Barker A. Absence of a functional *erm* gene in isolates of *Mycobacterium immunogenium* and the *Mycobacterium mucogenicum* group, based on in vitro clarithromycin susceptibility. *J Clin Microbiol.* 2015;53(3):875–8.
- Esteban J, Martín-de-Hijas NZ, Ortiz A, Kinnari TJ, Bodas Sánchez A, Gadea I, Fernández-Roblas R. Detection of *LfrA* and *tap* efflux pump genes among clinical isolates of non-pigmented rapidly growing mycobacteria. *Int J Antimicrob Agents.* 2009;34:454–6.

17. Ito T, Maruyama F, Sawai K, Nozaki K, Otsu K, Ohya K. Draft genome sequence of *Mycobacterium virginiense* strain GF75, isolated from the mud of a swine farm in Japan. *Genome Announc.* 2018;6:e00362–418.
18. Adékambi T, Colson P, Drancourt M. *rpoB*-based identification of non-pigmented and late-pigmenting rapidly growing mycobacteria. *J Clin Microbiol.* 2003;41:5699–708.
19. McNabb A, Eisler D, Adie K, Amos M, Rodrigues M, Stephens G, Black WA, Isaac-Renton J. Assessment of partial sequencing of the 65-kilodalton heat shock protein gene (*hsp65*) for routine identification of *Mycobacterium* species isolated from clinical sources. *J Clin Microbiol.* 2004;42:3000–11.
20. Nurk S, Bankevich A, Antipov D, Gurevich AA, Korobeynikov A, Lapidus A, Pribelski AD, Pyshkin A, Sirotkin A, Sirotkin Y, Stepanauskas R, Clingenpeel SR, Woyke T, McLean JS, Lasken R, Tesler G, Alekseyev MA, Pevzner PA. Assembling single-cell genomes and mini-metagenomes from chimeric MDA products. *J Comput Biol.* 2013;20:714–37.
21. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS ONE.* 2014;9:e112963.
22. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol.* 2017;13:e1005595.
23. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.* 2015;25:1043–55.
24. Menzel P, Ng KL, Krogh A. Fast and sensitive taxonomic classification for metagenomics with Kaiju. *Nat Commun.* 2016;7:11257.
25. Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML, Delmont TO. Anvi'o: an advanced analysis and visualization platform for omics data. *PeerJ.* 2015;3:e1319.
26. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res.* 2016;44:6614–24.
27. Richter M, Rosselló-Móra R, Glöckner FO, Peplies J. JSpeciesWS: a Web server for prokaryotic species circumscription based on pairwise genome comparison. *Bioinformatics.* 2016;32:929–31.
28. Ho YS, Adroub SA, Aleisa F, Mahmood H, Othoum G, Rashid F, Zaher M, Ali S, Bitter W, Pain A, Abdallah AM. Complete genome sequence of *Mycobacterium fortuitum* subsp. *fortuitum* type strain DSM46621. *J Bacteriol.* 2012;194:6337–8.
29. Pang S, Renvoisé A, Perret C, Guinier M, Chelghoum N, Brossier F, Capton E, Jarlier V, Sougakoff W. Whole-genome sequence of *Mycobacterium abscessus* clinical strain V06705. *Genome Announc.* 2013;1:e00690–713.
30. Asmar S, Rascovan N, Robert C, Drancourt M. Draft genome sequence of *Mycobacterium mucogenicum* strain CSUR P2099. *Genome Announc.* 2015;3:e01369–415.
31. Hasan NA, Davidson RM, de Moura VC, Garcia BJ, Reynolds PR, Epperson LE, Farias-Hesson E, DeGroot MA, Jackson M, Strong M. Draft genome sequence of *Mycobacterium chelonae* type strain ATCC 35752. *Genome Announc.* 2015;3:e00536–615.
32. Phelippeau M, Robert C, Croce O, Raoult D, Drancourt M. Draft genome sequence of *Mycobacterium neoaurum* strain DSM 44074T. *Genome Announc.* 2014;2:e00699–714.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

