

CORRECTION

Open Access



Correction: Differences in adoption of COVID-19 pandemic related preventive behaviour by viral load suppression status among people living with HIV during the first wave of the pandemic

Morenike Oluwatoyin Folayan^{1,2*}, Roberto Ariel Abeldaño Zuñiga^{1,3,4}, Nourhan M. Aly^{1,5}, Passent Ellakany^{1,6}, Ifeoma E. Idigbe^{1,7}, Mohammed Jafer^{1,8}, Folake B. Lawal^{1,9}, Zumama Khalid^{1,10}, Joanne Lusher^{1,11}, Jorma I. Virtanen^{1,12} and Annie L. Nguyen^{1,13}

Correction: BMC Research Notes (2023) 16:90
<https://doi.org/10.1186/s13104-023-06363-6>

Following the publication of the original article [1] it was brought to our attention that Table 1 contained some errors inadvertently introduced during typesetting.

The correct Table 1 is shown in this erratum:

The original article can be found online at <https://doi.org/10.1186/s13104-023-06363-6>.

*Correspondence:

Morenike Oluwatoyin Folayan
toyinukpong@yahoo.co.uk

¹ Mental Health and Wellness Study Group, Obafemi Awolowo University, Ile-Ife, Nigeria

² Department of Child Dental Health, Obafemi Awolowo University, Ile-Ife, Nigeria

³ Faculty of Social Sciences, University of Helsinki, Helsinki, Finland

⁴ Postgraduate Department, University of Sierra Sur, Oaxaca, Mexico

⁵ Department of Pediatric Dentistry and Dental Public Health, Faculty of Dentistry, Alexandria University, Alexandria, Egypt

⁶ Department of Substitutive Dental Sciences, College of Dentistry, Imam Abdulrahman Bin Faisal University, Dammam, Saudi Arabia

⁷ Clinical Sciences Department, Nigerian Institute of Medical Research, Lagos, Nigeria

⁸ Department of Preventive Dental Sciences, Faculty of Dentistry, Jazan University, Jazan, Saudi Arabia

⁹ Department of Periodontology and Community Dentistry, University of Ibadan and University College Hospital, Ibadan, Nigeria

¹⁰ Department of Health Sciences, University of Genova, Genoa, GE 16132, Italy

¹¹ Regent's University, London, UK

¹² Faculty of Medicine, University of Turku, Turku, Finland

¹³ Department of Family Medicine, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Table 1 Association between viral load, antiretroviral adherence profile and the adoption of COVID-19 pandemic preventive behaviour status among PLHIV during the first wave of the pandemic (N = 680)

Variables	Total	Physical distancing			Face mask			Handwashing			Work remotely		
		No	Yes	AOR; 95% CI; p value	No	Yes	AOR; 95% CI; p value	No	Yes	AOR; 95% CI; p value	No	Yes	AOR; 95% CI; p value
	N = 680	221 (32.5%)	459 (67.5%)		99 (14.6%)	581 (85.4%)		132 (19.4%)	548 (80.6%)		540 (79.4%)	140 (20.6)	
	n (%)	n (%)	n (%)		n (%)	n (%)		n (%)	n (%)		n (%)	n (%)	
Viral load													
Undetectable	488 (71.8)	157 (32.2)	331 (67.8)	1	56 (11.5)	432 (88.5)	1	86 (17.6)	402 (82.4)	1	389 (79.7)	99 (20.3)	1
Detectable	192 (28.2)	64 (33.3)	128 (66.7)	1.06; 0.72–1.55; p = 0.74	43 (22.4)	149 (77.6)	0.44; 0.28–0.69; p < 0.01	46 (24.0)	146 (76.0)	0.64; 0.42–0.97; p = 0.03	151 (78.6)	41 (21.4)	0.99; 0.64–1.53; p = 0.98
Adherence to ART													
No	151 (22.2)	53 (35.1)	98 (64.9)	1	23 (15.2)	128 (84.8)	1	30 (19.9)	121 (80.1)	1	107 (70.9)	44 (29.1)	1
Yes	529 (77.8)	168 (31.8)	361 (68.2)	1.04; 0.69–1.58; p = 0.83	76 (14.4)	453 (85.6)	0.99; 0.57–0.71; p = 0.98	102 (19.3)	427 (80.7)	1.11; 0.69–1.81; p = 0.65	433 (81.9)	96 (18.1)	0.60; 0.38–0.94; p = 0.02
Age	39.9 (11.3)	39.1 (11.6)	40.3 (11.2)	1.01; 1.00–1.03; p = 0.38	40.9 (11.3)	39.7 (11.3)	0.99; 0.97–1.00; p = 0.51	39.3 (10.1)	40.0 (11.6)	1.07; 0.98–1.02; p = 0.46	39.6 (11.6)	40.8 (10.3)	1.01; 0.99–1.03; p = 0.17
Sex													
Male	277 (40.8)	89 (32.1)	188 (67.9)	1	40 (14.4)	237 (85.6)	1	48 (17.3)	229 (82.7)	1	217 (78.3)	60 (21.7)	1
Female	395 (58.1)	129 (32.7)	266 (67.3)	0.97; 0.69–1.38; p = 0.89	56 (14.2)	339 (85.8)	0.94; 0.59–0.48; p = 0.80	82 (20.8)	313 (79.2)	0.81; 0.54–1.22; p = 0.32	316 (80.0)	79 (20.0)	0.93; 0.63–1.39; p = 0.75
Intersex	3 (0.4)	2 (66.7)	1 (33.3)	0.67; 0.04–9.71; p = 0.77	1 (33.3)	2 (66.7)	0.56; 0.03–0.87; p = 0.68	1 (33.3)	2 (66.7)	0.53; 0.03–7.87; p = 0.64	2 (66.7)	1 (33.3)	3.30; 0.22–49.46; p = 0.38
Decline to answer	5 (0.7)	1 (20.0)	4 (80.0)	1.72; 0.18–15.82; p = 0.62	2 (40.0)	3 (60.0)	0.20; 0.03–0.29; p = 0.09	1 (20.0)	4 (80.0)	0.96; 0.10–8.94; p = 0.97	5 (100.0)	0 (0.0)	Not calculated (p = 0.99)
Educational level													
No formal	24 (3.5)	20 (83.3)	4 (16.7)	0.66; 0.02–0.20; p < 0.01	8 (33.3)	16 (66.7)	0.30; 0.11–0.79; p = 0.01	6 (25.0)	18 (75.0)	0.49; 0.17–1.39; p = 0.18	21 (87.5)	3 (12.5)	0.24; 0.06–0.89; p = 0.03
Primary	57 (8.4)	30 (52.6)	27 (47.4)	0.34; 0.19–0.60; p < 0.01	14 (24.6)	43 (75.4)	0.43; 0.21–0.87; p = 0.01	12 (21.1)	45 (78.9)	0.78; 0.38–1.60; p = 0.50	53 (93.0)	4 (7.0)	0.19; 0.06–0.55; p < 0.01
Secondary	239 (35.1)	72 (30.1)	167 (69.9)	0.94; 0.65–1.36; p = 0.75	31 (13.0)	208 (87.0)	0.96; 0.58–1.59; p = 0.89	51 (21.3)	188 (78.7)	0.82; 0.53–1.26; p = 0.37	203 (84.9)	36 (15.1)	0.53; 0.34–0.82; p < 0.001

Table 1 (continued)

Variables	Total	Physical distancing		Face mask		Handwashing		Work remotely		AOR; 95% CI; p value
		No	Yes	No	Yes	No	Yes	No	Yes	
	N = 680	221	459	99	581	132	548	540	140	
		(32.5%)	(67.5%)	(14.6%)	(85.4%)	(19.4%)	(80.6%)	(79.4%)	(20.6)	
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	
Tertiary	360 (52.9)	99 (27.5)	261 (72.5)	46 (12.8)	314 (87.2)	63 (17.5)	297 (82.5)	263 (73.1)	97 (26.9)	1
Depression										
No	52 (81.2)	177 (32.1)	375 (67.9)	86 (15.6)	466 (84.4)	124 (22.5)	428 (77.5)	453 (82.1)	99 (17.9)	1
Yes	128 (18.8)	44 (34.4)	84 (65.6)	13 (10.2)	115 (89.8)	8 (6.3)	120 (93.7)	87 (68.0)	41 (32.0)	2.10; 1.33–3.31; p < 0.01
										4.80; 2.24–10.27; p < 0.01
										1.90; 0.99–3.68; p = 0.05
										1.09; 0.70–1.69; p = 0.70

ART antiretroviral therapy

The original article has been corrected.

Published online: 22 August 2023

Reference

1. Folayan MO, Zuñiga RAA, Aly NM, Ellakany P, Idigbe IE, Jafer M, Lawal FB, Khalid Z, Lusher J, Virtanen JI, Nguyen AL. Differences in adoption of COVID-19 pandemic related preventive behaviour by viral load suppression status among people living with HIV during the first wave of the pandemic. *BMC Res Notes*. 2023;16:90. <https://doi.org/10.1186/s13104-023-06363-6>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.