

RESEARCH NOTE

Open Access



Bacterial endotoxin lipopolysaccharides regulate gene expression in human colon cancer cells

Heping Cao^{1*}

Abstract

Objective Lipopolysaccharide (LPS) is a major cell wall component of gram-negative bacteria. Colon bacteria contribute to LPS which promotes colon cancer metastasis. The objective of this study was to survey the effect of LPS on cell viability and gene expression of 55 molecular targets in human colon cancer cells.

Results LPS did not affect the viability of COLO 225 cells under the culture conditions but affected the expression of a number of genes important in inflammatory responses and cancer development. LPS increased TTP family, GLUT family and DGAT1 mRNA levels but decreased DGAT2a and DGAT2b expression in the human colon cancer cells. LPS also increased COX2, CXCL1, ELK1, ICAM1, TNFSF10 and ZFAND5 but decreased BCL2L1, CYP19A1 and E2F1 mRNA levels in the colon cancer cells. These data suggest that LPS has profound effects on gene expression in human colon cancer cells.

Keywords Colon cancer cell, Cytotoxicity, Gene expression, Lipopolysaccharide

Introduction

Colon cancer is one of the deadliest diseases in the World. The risk of developing colorectal cancer is approximately 4.0% for men and women in 2021 during the lifetime (<https://www.cancer.org/cancer/types/colorectal-cancer/about/key-statistics.html>). It is urgently needed to fully understand the mechanism of developing colon cancer and explore ways to ease the burden of the healthcare crisis.

Lipopolysaccharide (LPS) is a major cell wall component of gram-negative bacteria. Intact LPS is made up of three structural components [1]: a hydrophobic lipid section; a hydrophilic core polysaccharide chain, and a

repeating hydrophilic O-antigenic oligosaccharide side chain. LPS is a heat-stable endotoxin which normally protects gram-negative bacteria against bile salts and lipophilic antibiotics.

LPS was proposed to have antitumor effect in several experimental models [2]. A number of studies explored the effect of LPS on gene expression in colon cancer cells, but they were focused on a few targets in the reported research. LPS induced TGF β and HGF production [3], promoted NF κ B (NFkappaB) activation [4] and increased the migratory capacity [5] in colon cancer cells.

The objective of this study was to survey the effect of LPS on cell viability and gene expression of 55 molecular targets in human colon cancer cells. The 55 molecular targets belong to several important pathways, whose expression is affected by the plant toxin gossypol in cancer cells [6–14] and macrophages [15, 16] or regulated by ZFP36/TTP in tumor cells [17–25] and macrophages [26, 27], as well as cinnamon polyphenol extract [28, 29] (Table 1). The results showed that LPS had minimal

*Correspondence:

Heping Cao
Heping.Cao@usda.gov

¹ United States Department of Agriculture, Agricultural Research Service, Southern Regional Research Center, 1100 Allen Toussaint Boulevard, New Orleans, LA 70124, USA



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Table 1 Basal mRNA level, reference mRNA and LPS effects on gene expression

ID	mRNA	Name	DMSO (n = 18)		LPS (n = 24)		LPS/DMSO
			Mean ± Std	Fold	Mean ± Std	Fold	Fold
H1	Ahrr1	Aryl hydrocarbon receptor	33.91 ± 1.22	0.05	37.63 ± 3.98	0.00	0.06
H2	Bcl2	B-cell lymphoma 2	29.68 ± 1.02	1.00	29.02 ± 1.16	1.00	1.00
H3	Bcl2l1	B-cell lymphoma 2 like 1	28.10 ± 2.27	2.99	27.72 ± 1.67	2.45	0.86
H4	Bnip3	BCL2 protein-interacting protein 3	27.94 ± 1.04	3.32	26.78 ± 1.50	4.72	1.30
H5	Cd36	Cluster of differentiation 36/fatty acid translocase	28.74 ± 1.25	1.92	27.65 ± 1.25	2.58	1.33
H6	Claudin1	Maintain tissue integrity and water retention	30.65 ± 5.90	0.51	31.23 ± 5.28	0.22	0.33
H7	Cox1	Cyclooxygenase 1	37.05 ± 5.88	0.01	39.19 ± 3.49	0.00	0.03
H8	Cox2	Cyclooxygenase 2	30.28 ± 1.33	0.66	31.79 ± 3.07	0.15	0.30
H9	Csnk2a1	Casein kinase 2 alpha 1	26.35 ± 2.10	10.05	26.12 ± 1.56	7.43	0.71
H10	Ctsb	Cathepsin B	28.47 ± 3.00	2.31	28.97 ± 2.31	1.03	0.53
H11	Cxcl1	Chemokine (C-X-C motif) ligand 1	32.81 ± 2.65	0.11	35.27 ± 5.69	0.01	0.18
H12	Cyclind1	Cyclin D1	34.88 ± 6.68	0.03	33.91 ± 4.96	0.03	2.59
H13	Cyp19a1	Cytochrome P450 family 19 subfamily A member 1	31.96 ± 3.39	0.21	28.56 ± 3.56	1.37	26.40
H14	Dgat1	Diacylglycerol O-acyltransferase 1	29.59 ± 1.93	1.06	30.17 ± 2.53	0.45	0.46
H15	Dgat2a	Diacylglycerol O-acyltransferase 2a	32.30 ± 2.13	0.16	33.30 ± 4.47	0.05	0.25
H16	Dgat2b	Diacylglycerol O-acyltransferase 2b	31.43 ± 1.71	0.30	31.19 ± 2.49	0.22	0.66
H17	E2f1	E2F transcription factor 1	29.82 ± 1.01	0.91	30.05 ± 2.75	0.49	0.57
H18	Elk1	ETS transcription factor	30.85 ± 2.79	0.44	31.73 ± 2.09	0.15	0.55
H19	Fas	Fas cell surface death receptor	31.60 ± 5.28	0.26	33.18 ± 4.49	0.06	0.26
H20	Gapdh	Glyceraldehyde-3-phosphate dehydrogenase	24.83 ± 4.17	28.71	25.07 ± 3.15	15.46	0.48
H21	Glut1	Glucose transporter 1	27.48 ± 2.72	4.57	29.21 ± 3.80	0.87	0.22
H22	Glut2	Glucose transporter 2	29.45 ± 2.00	1.17	29.05 ± 3.75	0.97	0.65
H23	Glut3	Glucose transporter 3	28.38 ± 1.21	2.45	27.78 ± 1.63	2.35	1.31
H24	Glut4	Glucose transporter 4	40.16 ± 5.08	0.00	41.70 ± 6.15	0.00	0.31
H25	Hif1a	Hypoxia inducible factor 1 subunit alpha	27.78 ± 2.29	3.72	27.67 ± 2.50	2.53	0.96
H26	Hmgr	3-Hydroxy-3-methylglutaryl-CoA reductase	27.85 ± 1.94	3.54	27.33 ± 1.25	3.22	0.94
H27	Hmox1	Heme oxygenase 1	30.11 ± 1.35	0.74	29.39 ± 1.23	0.77	0.96
H28	Hua	Human antigen a	32.98 ± 3.77	0.10	32.39 ± 3.71	0.10	0.75
H29	Icam1	Intercellular adhesion molecule 1/CD54	34.27 ± 4.65	0.04	36.49 ± 5.75	0.01	0.09
H30	Inos	Inducible nitric oxide synthase	ud	ud	ud	ud	ud
H31	Insr	Insulin receptor	29.99 ± 3.43	0.81	31.64 ± 3.23	0.16	0.29
H32	Il2	Interleukin 2	31.69 ± 1.08	0.25	30.31 ± 1.23	0.41	1.43
H33	Il6	Interleukin 6	29.51 ± 1.21	1.12	27.60 ± 1.22	2.67	1.95
H34	Il8	Interleukin 8	29.37 ± 1.08	1.24	28.92 ± 1.79	1.07	0.99
H35	Il10	Interleukin 10	36.16 ± 9.42	0.01	34.04 ± 11.21	0.03	1.10
H36	Il12	Interleukin 12	38.14 ± 3.63	0.00	32.53 ± 7.75	0.09	20.16
H37	Il16	Interleukin 16	28.45 ± 1.13	2.33	27.04 ± 4.88	3.94	1.73
H38	Il17	Interleukin 17	29.90 ± 1.30	0.85	28.92 ± 1.77	1.07	1.02
H39	Leptin	Body fat and obesity hormone	30.46 ± 5.47	0.58	28.98 ± 1.34	1.03	1.29
H40	Map1lc3a	Microtubule-associated proteins 1 light chain 3A	30.03 ± 1.82	0.78	29.03 ± 1.56	0.99	1.08
H41	Map1lc3b	Microtubule-associated proteins 1 light chain 3B	26.60 ± 1.64	8.44	26.93 ± 2.65	4.25	0.51
H42	Nfkb	Nuclear factor kappa B	31.25 ± 3.21	0.34	32.91 ± 5.03	0.07	0.28
H43	P53	Tumor suppressor	31.18 ± 2.46	0.35	30.71 ± 1.61	0.13	0.92
H44	Pim1	Proto-oncogene serine/threonine-protein kinase	29.42 ± 0.99	1.19	29.49 ± 1.57	0.72	0.59
H45	Pparr	Peroxisome proliferator-activated receptor gamma	29.36 ± 1.54	1.24	29.89 ± 1.36	0.54	0.62
H46	Rab24	Ras-related oncogene 24	41.98 ± 2.85	0.00	44.31 ± 5.63	0.00	0.15
H47	Rpl32	Ribosomal protein L32 (60S ribosomal unit)	24.59 ± 3.89	33.88	24.98 ± 3.10	16.40	0.55
H48	Tnf	Tumor necrosis factor	31.25 ± 1.76	0.34	30.44 ± 1.55	0.37	1.03

Table 1 (continued)

ID	mRNA	Name	DMSO (n = 18)		LPS (n = 24)		LPS/DMSO
			Mean ± Std	Fold	Mean ± Std	Fold	
H49	Tnfsf10	Tumor necrosis factor superfamily, member 10	28.24 ± 1.68	2.71	28.06 ± 1.41	1.94	0.83
H50	Ulk2	Unc-51 like autophagy activating kinase 2	29.54 ± 1.02	1.10	28.33 ± 1.36	1.61	1.27
H51	Vegf	Vascular endothelial growth factor	37.19 ± 6.99	<i>0.01</i>	37.26 ± 7.71	<i>0.00</i>	0.82
H52	Zfand5	Zinc finger AN1-type containing 5	27.47 ± 1.41	4.61	26.95 ± 1.50	4.20	1.00
H53	Zfp36	Zinc finger protein 36	29.04 ± 2.01	1.55	29.15 ± 1.85	0.91	0.69
H54	Zfp36l1	Zinc finger protein 36 like 1	29.78 ± 3.02	0.93	29.67 ± 2.70	<i>0.64</i>	0.80
H55	Zfp36l2	Zinc finger protein 36 like 2	41.81 ± 3.74	<i>0.00</i>	42.67 ± 2.47	<i>0.00</i>	0.37

The fold was calculated using the mean data. Bold with italics: Genes with mRNA levels at least twofold of Bcl2. Italics: Genes with mRNA levels less than 50% of Bcl2 undetected

effect on cell viability but had a profound effect on gene expression at the mRNA levels in the human colon cancer cells.

Main text

Methods

Human colon cancer cells (COLO 205) were maintained in RPMI-1640 medium (Gibco) containing 10% (v:v) fetal bovine serum, 0.1 million units/L penicillin, 100 mg/L streptomycin, and 2 mmol/L L-glutamine at 37 °C with 5% CO₂. Cancer cells (0.5 mL) were subcultured at 1 × 10⁵ cells/mL density and treated for 2 and 24 h with 0–1000 ng/mL of LPS extracted from *E. coli* serotype K235 and purified by gel filtration (Sigma, St. Louis, MO) (“0” treatment represents the control with 1% DMSO in all treatments). Cell cytotoxicity was determined by spectrophotometer at A570 nm using the MTT based-In Vitro Toxicology Assay Kit (Sigma) [30].

The effect of LPS on gene expression was evaluated by quantitative real-time PCR analysis (qPCR). Fifty-five genes were selected for qPCR analysis (Table 1). Human colon cancer cells in triplicate were treated with LPS for 8 h. RNA isolation and cDNA synthesis were performed as described [31]. The SYBR Green qPCR assays were described previously [32, 33]. BCL2 mRNA was selected as the internal reference based on our previously analysis [14] and date presented in the “Results” section. The $2^{-\Delta CT}$ and $2^{-\Delta\Delta CT}$ method of relative quantification was used to determine the fold change in expression [34]. The data represent the mean and standard deviation (the number of independent qPCR data ‘n’ is indicted in the tables and figure legends).

Results

Effect of LPS on cell viability

MTT method assessed cell cytotoxicity of human colon cancer cells (COLO 225) after the cells were treated with up to 1000 ng/mL of LPS for 2 and 24 h. MTT assay did

not show significant effect of LPS on the viability of the human colon cancer cells under the culture conditions (Data not shown).

Basal expression level

To provide a basis for gene expression comparison in the colon cancer cells, the relative mRNA levels of 55 genes were estimated by SYBR Green qPCR assay. The qPCR assay showed that the cycle of threshold (C_T) of BCL2 mRNA was 29.68 ± 1.02 (mean ± standard deviation, n = 18, means the calculation was performed from 18 independent qPCR data) (Table 1). GAPDH and RPL32 mRNA levels were the most abundant with 29- and 34-fold of BCL2 mRNA, respectively. INOS mRNA was undetectable. AHRR1, COX1, CYCLIND1, GLUT4, ICAM1, IL10, IL12, RAB24, VEGF and ZFP36L2 mRNA levels were detected with less than 5% of BCL2 mRNA in the colon cancer cells (Table 1).

Selection of reference mRNA

The ideal reference gene for qPCR assay is stably expressed under the experimental conditions. This can be estimated by the standard deviations among the treatments. The less of standard deviation of C_T among the LPS treatments indicates the more stable expression of the gene. The C_T value of BCL2 mRNA was 29.02 ± 1.16 (mean ± standard deviation, n = 24), one of the least varied mRNAs (Table 1). GAPDH and RPL32 mRNAs are widely used as references for qPCR assays in other mammalian cells such as adipocytes and macrophages [28, 29, 35, 36], but GAPDH and RPL32 mRNA levels had much larger standard deviations (ΔCT was 3.15 and 3.10, respectively) and the most abundantly expressed with 15.5- and 16.4-fold of BCL2 mRNA, respectively in the human colon cancer cells (Table 1). The large standard deviations and high expression levels of GAPDH and RPL32 mRNAs made them not good internal references for qPCR assays in the human colon

cancer cells. BCL2 was among the least regulated genes by LPS and therefore suitable as the internal reference for the qPCR analyses.

Effect of LPS on overall gene expression

To provide a general idea how these genes were expressed in the cultured colon cancer cells with or without LPS treatment, the pooled qPCR data were analyzed using BCL2 mRNA as the internal reference and DMSO treatment as the sample control. LPS upregulated the expression of three mRNAs with at least twofold of the control but decreased the expression of 16 mRNAs with less than 50% of the control. The up-regulated 3 mRNAs were CYCLIND1, CYP19A1 and IL12 (Table 1). The down-regulated 17 mRNAs were AHRR1, CLAUDIN1, COX1, COX2, CXCL1, DGAT1, DGAT2a, FAS, GAPDH, GLUT1, GLUT4, ICAM1, INSR, NFKB, RAB24 and ZFP36L2 (Table 1). However, it is worth mentioning that the expression patterns based on pooled data from various concentrations may not completely in agreement with those of the detailed analysis of pair-wised comparison between the treatment and DMSO control as detailed below.

Effect of LPS on gossypol-reported gene expression

Several genes were shown previously to be regulated by plant toxin gossypol in cancer cells and macrophages. Here, we analyzed the expression of the same group of genes including BNIP3, CYP19A1, FAS, HuA, P53, PPARR and TNFSF10 genes under various concentrations of LPS in the colon cancer cell line using BCL2 as the internal reference gene [31]. In general, this group of genes were expressed lower than BCL2 except BINP3 and TNFSF10 (Table 1). LPS increased mRNA levels of P53, PPARR and TNFSF10 genes but decreased that of CYP19A1 gene (Fig. 1A). The effects of bacterial endotoxin LPS on the expression of this group of genes were different from those of the plant toxin gossypol which inhibited the expression of all these genes except PPARR gene to a large extent in the same human colon cancer cells [31].

Effect of LPS on DGAT gene expression

Diacylglycerol acyltransferases (DGATs) catalyze the rate-limiting step of triacylglycerol biosynthesis by esterifying *sn*-1,2-diacylglycerol with a long-chain fatty acyl-CoA. DGAT2 mRNA is the major DGAT mRNA in mouse adipocytes and macrophages [33, 37], but DGAT1 mRNA was the major form in the human colon cancer cells (Table 1). LPS treatment under higher concentration increased DGAT1 mRNA levels but decreased DGAT2a and DGAT2b expression in the human colon cancer cells (Fig. 1B).




Fig. 1 Effect of LPS on the expression of gossypol-regulated genes as well as DGAT and GLUT family genes

Effect of LPS on GLUT gene expression

Glucose transporter (GLUT) family proteins are responsible for glucose uptake in mammalian cells. GLUT1 mRNA was the major form of GLUT mRNAs but GLUT4 mRNA was barely detected in the colon cancer cells (Table 1). LPS treatment significantly increased GLUT2 and GLUT3 mRNA levels but only high concentration of LPS at 1000 ng/mL increased GLUT1 mRNA level (Fig. 1C).

Effect of LPS on TTP and TTP-mediated gene expression

Tristetraprolin (TTP/ZFP36) family proteins control the mRNA stability of some cytokines [38]. qPCR showed that TTP and ZFP36L1 genes were expressed in similar levels but ZFP36L2 mRNA was barely detectable in the colon cancer cells (Table 1). ZFP36, ZFP36L1 and ZFP36L2 mRNA levels were generally increased in the colon cancer cells by high concentration of LPS treatment (Fig. 2A).




Fig. 2 Effect of LPS on TTP and IL, proinflammatory and other family gene expression

Effect of LPS on proinflammatory gene expression

TTP family proteins down-regulate the stability of several proinflammatory cytokine and enzyme mRNAs including those coding for tumor necrosis factor-alpha (TNF α) [39], granulocyte–macrophage colony-stimulating factor/colony-stimulating factor 2 (GM-CSF/CSF2) [40] and cyclooxygenase 2/prostaglandin-endoperoxide synthase 2 (COX2/PTGS2) [24]. qPCR showed that all the tested proinflammatory mRNAs except TNFSF10 mRNA were expressed much lower than that of TTP in the colon cancer cells (Table 1). LPS increased COX2 and TNFSF10 mRNA levels but did not exhibit significant effect on HuA, LEPTIN and TNF mRNA levels in the human colon cancer cells (Fig. 2B). COX1 and VEGF mRNA levels were too low to be reliable (Table 1).

Effect of LPS on IL gene expression

TTP family proteins also regulate the stability of several interleukin (IL) mRNAs coding for IL2 [41], IL6 [42], IL8 [43], IL10 [44], IL12 [45], IL16 [23] and IL17 [46]. SYBR Green qPCR showed that IL10 and IL12 mRNAs were barely expressed and IL2 mRNA was low, whereas the other ILs were expressed in similar levels to TTP, which were several fold higher than IL2 mRNA in the human colon cancer cells (Table 1). The qPCR assays showed that LPS did not have significant effect on IL mRNA levels in the colon cancer cells (Fig. 2C).

Effect of LPS on TTP-targeted other gene expression

A number of other TTP-mediated mRNAs have been reported in the literature. The basal levels of some mRNAs were higher than that of TTP mRNA (BCL2L1, CSNK2A1, HIF1a and ZFAND5) but the others were lower than that of TTP mRNA (AHRR1, CXCL1, E2F1, ELK1, HMOX1 and ICAM1) (Table 1). qPCR showed that LPS increased CXCL1, ELK1, ICAM1 and ZFAND5 mRNA levels, but decreased BCL2L1 and E2F1 mRNA levels in the colon cancer cells (Fig. 2D).

Discussion

Colon bacteria contribute to a large quantity of LPS which could promote colon cancer metastasis. In this study, we surveyed the effect of LPS on cell viability and expression of 55 genes at the mRNA levels in human colon cancer cells. The data confirmed that BCL2 was the most stable mRNA among the 55 mRNAs and suitable as the reference mRNA for qPCR analyses in human colon cancer cells [31]. We observed that LPS did not affect the viability of the cells but affected the expression of a number of genes important in inflammatory responses and cancer development under the culture conditions.

The following findings are worthy of discussion. (1) High concentration of LPS increased TTP family gene expression in the human colon cancer cells, in agreement with the previous results using mouse macrophages [29, 47]. (2) LPS increased GLUT1, GLUT2 and GLUT3

mRNA levels in the human colon cancer cells, suggesting that LPS maybe able to increase glucose transport into the cancer cells since GLUT family proteins are responsible for glucose uptake in mammalian cells [27, 33]. (3) LPS treatment under higher concentration increased DGAT1 mRNA levels (the major form of DGATs) but decreased DGAT2a and DGAT2b expression in the human colon cancer cells, suggesting that LPS has limited effect on triacylglycerol biosynthesis in the colon cancer cells. (4) LPS increased COX2 mRNA levels in this study, in contrast to a previous study [48], which might be due to the cell type (COLO 225 vs. Coco-2) and/or detection methods (qPCR vs. western blot) used in the two studies. (5) LPS did not show any significant effect on HIF1a gene expression in COLO 225 cells, similar to those using MC-38 mouse colon cancer cells [49]. (6) LPS did not have significant effect on IL gene expression in this study, similar to those showing that LPS does not increase IL6, IL8 and IL15 expression in two human colon cancer cell lines [3], but differ from two reports about LPS effect on IL6 and IL8 mRNA levels in HT-29 cells [50, 51].

Limitations

A few limitations of this study are worthy of mentioning. First, the data were generated from one colon cancer cell line (COLO 225). It could be valuable to expand the research with other cancer cell lines. Second, the dosage effect of LPS on mRNA levels was not strong and the standard deviations were large in some assays probably due to extremely sensitive qPCR assays. Third, it could be great to confirm mRNA data at the protein level. Finally, there is no functional analysis of LPS on intermediate steps between mRNA changes and cell viability. It is author's aim to present initial observations rather than in-depth study in this manuscript. Hopefully, more detailed studies could be performed when more resources are available for this type of study.

Acknowledgements

The author would like to express great appreciation to Dr. Kandan Sethumadhavan for technical assistance. The preliminary work of this study was presented at the 2022 American Chemical Society National Meeting in Chicago, IL [52].

Author contributions

HC designed and performed the experiments, analyzed the data and wrote the manuscript.

Funding

This work was supported by the USDA-ARS Quality and Utilization of Agricultural Products National Program 306 through ARS Research Projects 6054-41000-103-00-D and 6054-41000-113-00-D. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture. USDA is an equal opportunity provider and employer. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Availability of data and materials

The datasets generated during the current study are available in the NIH Gene Expression Omnibus (GEO) Database, accession number GSE200980 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE200980>). Materials are available from the author upon request.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The author declares no competing interests.

Received: 27 September 2022 Accepted: 6 September 2023

Published online: 13 September 2023

References

- Rietschel ET, Kirikae T, Schade FU, Mamat U, Schmidt G, Loppnow H, Ulmer AJ, Zähringer U, Seydel U, Di Padova F, et al. Bacterial endotoxin: molecular relationships of structure to activity and function. *FASEB J*. 1994;8(2):217–25.
- Lejeune P, Reisser D, Onier N, Lagadec P, Lindley I, Jeannin JF. Interleukin-8 has antitumor effects in the rat which are not associated with polymorphonuclear leukocyte cytotoxicity. *Cancer Immunol Immunother*. 1994;38(3):167–70.
- Yoshioka T, Morimoto Y, Iwagaki H, Itoh H, Saito S, Kobayashi N, Yagi T, Tanaka N. Bacterial lipopolysaccharide induces transforming growth factor beta and hepatocyte growth factor through toll-like receptor 2 in cultured human colon cancer cells. *J Int Med Res*. 2001;29(5):409–20.
- Ikebe M, Kitaura Y, Nakamura M, Tanaka H, Yamasaki A, Nagai S, Wada J, Yanai K, Koga K, Sato N, et al. Lipopolysaccharide (LPS) increases the invasive ability of pancreatic cancer cells through the TLR4/MyD88 signaling pathway. *J Surg Oncol*. 2009;100(8):725–31.
- Liu WT, Jing YY, Yan F, Han ZP, Lai FB, Zeng JX, Yu GF, Fan QM, Li R, Zhao QD, et al. LPS-induced CXCR4-dependent migratory properties and a mesenchymal-like phenotype of colorectal cancer cells. *Cell Adh Migr*. 2017;11(1):13–23.
- Yuan Y, Tang AJ, Castoreno AB, Kuo SY, Wang Q, Kuballa P, Xavier R, Shamji AF, Schreiber SL, Wagner BK. Gossypol and an HMT G9a inhibitor act in synergy to induce cell death in pancreatic cancer cells. *Cell Death Dis*. 2013;4: e690.
- Barba-Barajas M, Hernandez-Flores G, Lerma-Diaz JM, Ortiz-Lazareno PC, Dominguez-Rodriguez JR, de Barba-Barajas LCR, Jave-Suarez LF, Aguilar-Lemarroy AC, Guevara-Barraza MG, et al. Gossypol induced apoptosis of polymorphonuclear leukocytes and monocytes: involvement of mitochondrial pathway and reactive oxygen species. *Immunopharmacol Immunotoxicol*. 2009;31(2):320–30.
- Chang JS, Hsu YL, Kuo PL, Chiang LC, Lin CC. Upregulation of Fas/ Fas ligand-mediated apoptosis by gossypol in an immortalized human alveolar lung cancer cell line. *Clin Exp Pharmacol Physiol*. 2004;31(10):716–22.
- Dong Y, Mao B, Li L, Guan H, Su Y, Li X, Lian Q, Huang P, Ge RS. Gossypol enantiomers potently inhibit human placental 3beta-hydroxysteroid dehydrogenase 1 and aromatase activities. *Fitoterapia*. 2015;109:132–7.
- Huang YW, Wang LS, Chang HL, Ye W, Dowd MK, Wan PJ, Lin YC. Molecular mechanisms of (-)-gossypol-induced apoptosis in human prostate cancer cells. *Anticancer Res*. 2006;26(3A):1925–33.
- Kitada S, Kress CL, Krajewska M, Jia L, Pellicchia M, Reed JC. Bcl-2 antagonist apogossypol (NSC736630) displays single-agent activity in Bcl-2-transgenic mice and has superior efficacy with less toxicity compared with gossypol (NSC19048). *Blood*. 2008;111(6):3211–9.
- Ligueros M, Jeoung D, Tang B, Hochhauser D, Reidenberg MM, Sonenberg M. Gossypol inhibition of mitosis, cyclin D1 and Rb protein in human

- mammary cancer cells and cyclin-D1 transfected human fibrosarcoma cells. *BrJ Cancer*. 1997;76(1):21–8.
13. Yeow WS, Baras A, Chua A, Nguyen DM, Sehgal SS, Schrupp DS, Nguyen DM. Gossypol, a phytochemical with BH3-mimetic property, sensitizes cultured thoracic cancer cells to Apo2 ligand/tumor necrosis factor-related apoptosis-inducing ligand. *J Thorac Cardiovasc Surg*. 2006;132(6):1356–62.
 14. Cao H, Sethumadhavan K. Identification of Bcl2 as a stably expressed qPCR reference gene for human colon cancer cells treated with cottonseed-derived gossypol and bioactive extracts and bacteria-derived lipopolysaccharides. *Molecules*. 2022;27(21):7560.
 15. Cao H, Sethumadhavan K. Gossypol but not cottonseed extracts or lipopolysaccharides stimulates HuR gene expression in mouse cells. *J Funct Foods*. 2019;59:25–9.
 16. Cao H, Sethumadhavan K. Plant polyphenol gossypol induced cell death and its association with gene expression in mouse macrophages. *Biomolecules*. 2023;13(4):624.
 17. Essafi-Benkhadir K, Onesto C, Stebe E, Moroni C, Pages G. Tristetraprolin inhibits ras-dependent tumor vascularization by inducing VEGF mRNA degradation. *Mol Biol Cell*. 2007;18(11):4648–58.
 18. Florkowska M, Tymoszyk P, Balwierz A, Skucha A, Kochan J, Wawro M, Stalinska K, Kasza A. EGF activates TTP expression by activation of ELK-1 and EGR-1 transcription factors. *BMC Mol Biol*. 2012;13:8.
 19. Kim HK, Kim CW, Vo MT, Lee HH, Lee JY, Yoon NA, Lee CY, Moon CH, Min YJ, Park JW, et al. Expression of proviral integration site for Moloney murine leukemia virus 1 (Pim-1) is post-transcriptionally regulated by tristetraprolin in cancer cells. *JBiolChem*. 2012;287(34):28770–8.
 20. Lee HH, Lee SR, Leem SH. Tristetraprolin regulates prostate cancer cell growth through suppression of E2F1. *J Microbiol Biotechnol*. 2014;24(2):287–94.
 21. Lee HH, Kim WT, Kim DH, Park JW, Kang TH, Chung JW, Leem SH. Tristetraprolin suppresses AHR expression through mRNA destabilization. *FEBS Lett*. 2013;587(10):1518–23.
 22. Lee WH, Lee HH, Vo MT, Kim HJ, Ko MS, Im YC, Min YJ, Lee BJ, Cho WJ, Park JW. Casein kinase 2 regulates the mRNA-destabilizing activity of tristetraprolin. *J Biol Chem*. 2011;286(24):21577–87.
 23. Milke L, Schulz K, Weigert A, Sha W, Schmid T, Brune B. Depletion of tristetraprolin in breast cancer cells increases interleukin-16 expression and promotes tumor infiltration with monocytes/macrophages. *Carcinogenesis*. 2013;34(4):850–7.
 24. Sawaoka H, Dixon DA, Oates JA, Boutaud O. Tristetraprolin binds to the 3'-untranslated region of cyclooxygenase-2 mRNA. A polyadenylation variant in a cancer cell line lacks the binding site. *J Biol Chem*. 2003;278(16):13928–35.
 25. Sharma A, Bhat AA, Krishnan M, Singh AB, Dhawan P. Trichostatin-A modulates claudin-1 mRNA stability through the modulation of Hu antigen R and tristetraprolin in colon cancer cells. *Carcinogenesis*. 2013;34(11):2610–21.
 26. Cao H, Polansky MM, Anderson RA. Cinnamon extract and polyphenols affect the expression of tristetraprolin, insulin receptor, and glucose transporter 4 in mouse 3T3-L1 adipocytes. *Arch Biochem Biophys*. 2007;459(2):214–22.
 27. Cao H, Urban JF Jr, Anderson RA. Cinnamon polyphenol extract affects immune responses by regulating anti- and proinflammatory and glucose transporter gene expression in mouse macrophages. *J Nutr*. 2008;138(5):833–40.
 28. Cao H, Sethumadhavan K, Li K, Boue SM, Anderson RA. Cinnamon polyphenol extract and insulin regulate diacylglycerol acyltransferase gene expression in mouse adipocytes and macrophages. *Plant Foods Hum Nutr*. 2019;74(1):115–21.
 29. Cao H, Urban JF Jr, Anderson RA. Cinnamon polyphenol extract affects immune responses by regulating anti- and proinflammatory and glucose transporter gene expression in mouse macrophages. *J Nutr*. 2008;138(5):833–40.
 30. Cao H, Sethumadhavan K, Bland JM. Isolation of cottonseed extracts that affect human cancer cell growth. *SciRep*. 2018;8(1):10458.
 31. Cao H, Sethumadhavan K, Cao F, Wang TTY. Gossypol decreased cell viability and down-regulated the expression of a number of genes in human colon cancer cells. *Sci Rep*. 2021;11(1):5922.
 32. Cao H, Shockey JM. Comparison of TaqMan and SYBR Green qPCR methods for quantitative gene expression in tung tree tissues. *J Agric Food Chem*. 2012;60(50):12296–303.
 33. Cao H, Cao F, Roussel AM, Anderson RA. Quantitative PCR for glucose transporter and tristetraprolin family gene expression in cultured mouse adipocytes and macrophages. *In Vitro Cell Dev Biol Anim*. 2013;49(10):759–70.
 34. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods*. 2001;25(4):402–8.
 35. Cao H, Graves DJ, Anderson RA. Cinnamon extract regulates glucose transporter and insulin-signaling gene expression in mouse adipocytes. *Phytomedicine*. 2010;17(13):1027–32.
 36. Cao H, Urban JF Jr, Anderson RA. Insulin increases tristetraprolin and decreases VEGF gene expression in mouse 3T3-L1 adipocytes. *Obesity*. 2008;16(6):1208–18.
 37. Cao H. Identification of the major diacylglycerol acyltransferase mRNA in mouse adipocytes and macrophages. *BMC Biochem*. 2018;19(11):1–11.
 38. Blackshear PJ. Tristetraprolin and other CCHC tandem zinc-finger proteins in the regulation of mRNA turnover. *Biochem Soc Trans*. 2002;30(6):945–52.
 39. Carballo E, Lai WS, Blackshear PJ. Feedback inhibition of macrophage tumor necrosis factor-alpha production by tristetraprolin. *Science*. 1998;281(5379):1001–5.
 40. Carballo E, Lai WS, Blackshear PJ. Evidence that tristetraprolin is a physiological regulator of granulocyte-macrophage colony-stimulating factor messenger RNA deadenylation and stability. *Blood*. 2000;95(6):1891–9.
 41. Ogilvie RL, Abelson M, Hau HH, Vlasova I, Blackshear PJ, Bohjanen PR. Tristetraprolin down-regulates IL-2 gene expression through AU-rich element-mediated mRNA decay. *J Immunol*. 2005;174(2):953–61.
 42. Hochdorfer T, Tiedje C, Stumpo DJ, Blackshear PJ, Gaestel M, Huber M. LPS-induced production of TNF-alpha and IL-6 in mast cells is dependent on p38 but independent of TTP. *Cell Signal*. 2013;25(6):1339–47.
 43. Balakathiresan NS, Bhattacharyya S, Gutti U, Long RP, Jozwik C, Huang W, Srivastava M, Pollard HB, Biswas R. Tristetraprolin regulates IL-8 mRNA stability in cystic fibrosis lung epithelial cells. *Am J Physiol Lung Cell Mol Physiol*. 2009;296(6):L1012–8.
 44. Gaba A, Grivennikov SI, Do MV, Stumpo DJ, Blackshear PJ, Karin M. Cutting edge: IL-10-mediated tristetraprolin induction is part of a feedback loop that controls macrophage STAT3 activation and cytokine production. *J Immunol*. 2012;189(5):2089–93.
 45. Gu L, Ning H, Qian X, Huang Q, Hou R, Almourani R, Fu M, Blackshear PJ, Liu J. Suppression of IL-12 production by tristetraprolin through blocking NF-kappaB nuclear translocation. *J Immunol*. 2013;191(7):3922–30.
 46. Datta S, Novotny M, Pavicic PG Jr, Zhao C, Herjan T, Hartupej J, Hamilton T. IL-17 regulates CXCL1 mRNA stability via an AUUUA/tristetraprolin-independent sequence. *J Immunol*. 2010;184(3):1484–91.
 47. Cao H, Tuttle JS, Blackshear PJ. Immunological characterization of tristetraprolin as a low abundance, inducible, stable cytosolic protein. *JBiolChem*. 2004;279(20):21489–99.
 48. Arbabi S, Rosengart MR, Garcia I, Jelacic S, Maier RV. Epithelial cyclooxygenase-2 expression: a model for pathogenesis of colon cancer. *J SurgRes*. 2001;97(1):60–4.
 49. Muller-Edenborn K, Leger K, Glaus Garzon JF, Oertli C, Mirsaidi A, Richards PJ, Rehrauer H, Spielmann P, Hoogewijs D, Borsig L, et al. Hypoxia attenuates the proinflammatory response in colon cancer cells by regulating IkappaB. *Oncotarget*. 2015;6(24):20288–301.
 50. Qi HM, Zhu TM, Wang J. Regulation of immune suppressive cytokines by TLR4 activation in colon cancer cells. *Zhonghua Wei Chang Wai Ke Za Zhi*. 2009;12(4):413–5.
 51. Tang X, Zhu Y. TLR4 signaling promotes immune escape of human colon cancer cells by inducing immunosuppressive cytokines and apoptosis resistance. *OncolRes*. 2012;20(1):15–24.
 52. Cao H. Bacterial endotoxin lipopolysaccharides regulate gene expression in human colon cancer cells. In: American Chemical Society National Meeting: 08/23/2022 2022; McCormick Place Convention Center, Chicago, IL: American Chemical Society; 2022: PAPER ID: 3739772.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.