

SHORT REPORT

Open Access

A hyperactive *sleeping beauty* transposase enhances transgenesis in zebrafish embryos

Morgan Newman*, Michael Lardelli

Abstract

Background: Transposons are useful molecular tools for transgenesis. The 'sleeping beauty' transposon is a synthetic member of the Tc1/mariner transposon family. Davidson *et al.* (2003) previously described a vector for zebrafish transgenesis consisting of the inverted repeats of 'sleeping beauty' flanking the gene to be transposed. Subsequently, there have been attempts to enhance the transpositional activity of 'sleeping beauty' by increasing the activity of its transposase. Recently, Mates *et al.* (2009) generated a hyperactive transposase giving a 100-fold increased transposition rate in mouse embryos.

Findings: The aim of this experiment was to determine whether this novel hyperactive transposase enhances transgenesis in zebrafish embryos. Using our previously characterised *mitfa*-amyloid β -GFP transgene, we observed an eight-fold enhancement in transient transgenesis following detection of transgene expression in melanophores by whole mount *in-situ* hybridisation. However, high rates of defective embryogenesis were also observed.

Conclusion: The novel hyperactive 'sleeping beauty' transposase enhances the rate of transgenesis in zebrafish embryos.

Findings

Transposons direct integrations of single copies of genetic material into chromosomes [1] and are useful molecular tools for transgenesis in vertebrate species. They function by delivering a gene of interest to the chromosome in a cut and paste manner. The 'sleeping beauty' transposon is a synthetic member of the Tc1/mariner transposon family. The transposon was engineered from a consensus sequence of inactive fossil transposon sequences from various Salmonid fish genomes [2]. Sleeping beauty consists of the transposase gene flanked by terminal inverted repeats of direct repeats. The transposase protein catalyses the excision and integration of donor DNA into a TA dinucleotide site of a recipient genome [1]. The derived sleeping beauty vector system (SBT) has been shown to enhance production of transgenic animals in comparison to simple methods of transgenesis such as injection of naked DNA [3,4]. It is active in various vertebrate species such as fish, frogs, mice and rats [3,5-7]. There have been attempts to enhance the transpositional activity of the SBT, specifically by increasing the activity of the transposase. Almost every amino acid has

been changed to derive hyperactive mutants of the SB transposase and this has yielded modest increases in transpositional activity [8-11]. Recently, Mates *et al.* [5], used a large-scale genetic screen in mammalian cells to generate a hyperactive transposase that gave a ~100-fold enhancement of transpositional activity over the original SB transposase in mouse embryos.

Alzheimer's disease may be caused by the accumulation of amyloid β peptides in the brain [12]. Recently, we used the SBT system to generate a zebrafish melanophore model of amyloid β toxicity [7]. We generated transgenic zebrafish possessing human amyloid β under the control of the *mitfa* promoter that drives expression specifically in melanophores (dark pigment cells) using our vector pT2-*mitfa*-amyloid β -GFP. In that study the transposase mRNA was generated from the plasmid pSBRNAX that includes sequence from the 3' UTR of the *Xenopus* β -globin gene for mRNA stabilisation. In this experiment, we compared the rates of transient transgenesis in zebrafish embryos using the original transposase mRNA (SB10, generated from pSBRNAX [3]) or the hyperactive transposase mRNA (SB100, generated from pCMV(CAT)T7-SB100X [5]). It is important to note that the pCMV(CAT)T7-SB100X vector does not contain the *Xenopus*

* Correspondence: morgan.newman@adelaide.edu.au
Discipline of Genetics, The School of Molecular and Biomedical Science, The University of Adelaide, Adelaide, SA, 5005, Australia

β -globin 3' UTR sequences for mRNA stabilisation. Therefore, SB100 mRNA may not be as stable as SB10 mRNA and, once injected into the zebrafish embryos, may possibly degrade at a faster rate.

Zebrafish embryos were injected at the 1-cell stage with ~3 nl of linearised pT2-*mitfa*-amyloid β -GFP mixed with either SB100 or SB10 transposase mRNA (final concentration of DNA and mRNA is 25 ng/ μ l each) [6]. Embryos were permitted to develop until ~24 hour post-fertilisation (hpf) at which time their chorions were removed and they were fixed in 4% formaldehyde in a phosphate buffered saline solution. There was some variability in the normal development of individual embryos injected with the SB100 mRNA. From a total of 58 injected embryos, 22 showed developmental defects. Specifically, 6 showed defects in epiboly (but continued to develop later stage tissues) and 16 had trunk/somitogenesis development defects (Table 1).

Whole-mount *in-situ* transcript hybridization (WISH) was then performed on fixed embryos essentially as described by Jowett [13]. Since the GFP coding sequences in the pT2-*mitfa*-amyloid β -GFP transgene are transcribed and not translated a digoxigenin-labelled antisense EGFP cRNA probe was used, as previously described [7], to detect cells transcribing GFP (in general, this can also provide more sensitive detection of gene expression than observation of GFP fluorescence). Fixed embryos were stained for ~18 hours at 4°C, followed by ~6 hours at room temperature to be confident that all putative melanophores expressing the GFP transcript were revealed. There was some variability in the number of putative melanophores expressing the GFP transcript in individual embryos. However, of the 58 SB100 mRNA injected embryos, 14 (24%) (Figure 1F) had putative melanophores expressing the GFP transcript (see figure 1A-E) and of the 61 SB10 mRNA injected embryos, only 2 (3%) (Figure 1F) had putative melanophores expressing the GFP transcript. Therefore, injection of the SB100 mRNA resulted in an 8-fold enhancement of transient transgenesis in zebrafish

embryos. Interestingly, 10 out of 14, SB100 mRNA injected embryos with GFP transcript expression, also had the above mentioned trunk/somitogenesis development defects. This is consistent with higher rates of transgenesis being associated with higher rates of deformity [14]. To determine whether injection of the SB100 mRNA by itself might cause developmental defects, we determined the relative rates of defective embryos from injections of only linearised pT2-*mitfa*-amyloid β -GFP DNA, SB10 mRNA, SB100 mRNA or another transposase mRNA transcribed from the pCS-TP plasmid [15] (TOL2 mRNA) at 25 ng/ μ l. The results in Table 1 clearly show that only the SB100 mRNA causes increased developmental defects, indicating that the SB100 mRNA and not the transgene causes this. Embryos with epiboly defects are arrested in development at a stage before differentiation of melanophores is expected. Therefore, it is not possible to observe melanophore-specific GFP expression in these embryos. However, the possibility exists that these embryos also possess the transgene. Testing of the effects of a range of SB100 mRNA injection concentrations will be necessary to determine which concentration gives the optimum balance between transgenesis and embryo defect rates.

In their tests of SB100-driven transgenesis in fertilised mouse oocytes, Mates *et al.* (2009) did not observe a decreased survival rate relative to uninjected controls at day 7 of mouse embryogenesis and high rates of transgenesis were observed in mouse litters. However, the slower rate of cell division that occurs in cleavage stage mouse embryos relative to zebrafish embryos may mean that the transposase mRNA breaks down in the mouse zygotes before it can cause developmental defects.

The enhancement of transgenesis in zebrafish embryos from use of the novel hyperactive transposase was not ~100-fold greater than the transgenesis rate using the original SB transposase. Nevertheless, the observed 8-fold increase is a considerable improvement for two reasons. First, the amyloid β -GFP transgene is under the control of a tissue-specific promoter, *mitfa*, which directs expression of the transgene to melanophores. Melanophores make up only a small fraction of the total cells in a zebrafish embryo at 24 hpf. Thus, transient transgenesis is not expected to label this cell type frequently. Secondly, the SB10 mRNA is generated from pSBRNAX which has the *Xenopus* β -globin 3' UTR sequence for increased mRNA stability while the SB100 mRNA does not include such sequences. Therefore, the SB100 mRNA would be expected to degrade at a faster rate which might also affect transgenesis efficiency. If one considers that the germline transmission frequency of *mitfa*-amyloid β -GFP in the original study using SB10 mRNA was 20% (for a <3% rate of observable transient transgenesis), then the 8-fold enhancement of transient transgenesis observed in this study would presumably

Table 1 Results of injections of transposase mRNAs alone or with the *mitfa*-amyloid β -GFP transgene

Injection	Normal % (n)	Mild % (n)	Severe % (n)
Uninjected	98 (59)	2 (1)	0
<i>mitfa</i> -amyloid β -GFP only	94 (60)	3 (2)	3 (2)
-with SB100 mRNA	62 (36)	28 (16)	10 (6)
-with SB10 mRNA	97 (59)	3 (2)	0
SB100 mRNA only	40 (20)	16 (8)	44 (22)
SB10 mRNA only	83 (39)	8.5 (4)	8.5 (4)
TOL2 mRNA only	83 (38)	8.5 (4)	8.5 (4)

Percentage of embryos at 24 hpf with the number of embryos observed in parentheses.

Normal: wild type appearance. Mild: trunk/somitogenesis-like defects. Severe: epiboly-like defects.




Figure 1 Transgene expression after use of SB100x transposase. Embryos injected with transgene pT2-*mitfa*-amyloid β -GFP and SB100x mRNA were examined at 24 hpf for transgene expression by whole mount *in situ* transcript hybridisation against GFP sequences included in the transcript. A-D, lateral views of trunk region of embryos. E, dorsolateral view of hindbrain/spinal cord region of embryo. White arrowheads indicate some of the cells apparently expressing the transgene in a pattern consistent with expression in future melanophores in which the *mitfa* promoter is active. Size bars indicate approximately 100 μ m. Deformities caused by the SB100 transposase mean that some structures (e.g. the yolk extension) are an inconsistent size. F, Histogram showing percentage rates of transient transgenesis driven by SB10 and SB100 transposases.

further improve the rate of germline transgenesis in zebrafish. Overall, we conclude that the novel hyperactive 'sleeping beauty' transposase enhances the rate of transgenesis in zebrafish embryos.

Acknowledgements

Would like to thank Dr. Zsuzsanna Izsvak for providing the pCMV(CAT)T7-SB100X vector. This work was carried out under the auspices of the Animal Ethics Committee of The University of Adelaide. Research was supported by funds from the School of Molecular and Biomedical Research of The University of Adelaide, National Health and Medical research Council (NHMRC) Project Grant 453622 and the Cancer Council of South Australia.

Authors' contributions

MN completed experiments, participated in the design of the study and data analysis and drafted the manuscript. ML predominantly designed the study, participated in data analysis and revisions of the manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 21 June 2010 Accepted: 4 November 2010
Published: 4 November 2010

References

1. Wadman SA, Clark KJ, Hackett PB: Fishing for answers with transposons. *Mar Biotechnol (NY)* 2005, **7**:135-141.
2. Ivics Z, Hackett PB, Plasterk RH, Izsvak Z: Molecular reconstruction of Sleeping Beauty, a Tc1-like transposon from fish, and its transposition in human cells. *Cell* 1997, **91**:501-510.
3. Davidson AE, Balciunas D, Mohn D, Shaffer J, Hermanson S, Sivasubbu S, Cliff MP, Hackett PB, Ekker SC: Efficient gene delivery and gene expression in zebrafish using the Sleeping Beauty transposon. *Dev Biol* 2003, **263**:191-202.
4. Dupuy AJ, Clark K, Carlson CM, Fritz S, Davidson AE, Markley KM, Finley K, Fletcher CF, Ekker SC, Hackett PB, Horn S, Largaespada DA: Mammalian germ-line transgenesis by transposition. *Proc Natl Acad Sci USA* 2002, **99**:4495-4499.
5. Mates L, Chuah MK, Belay E, Jerchow B, Manoj N, Acosta-Sanchez A, Grzela DP, Schmitt A, Becker K, Matrai J, Ma L, Samara-Kuko E, Gysemans C, Pryputniewicz D, Miskey C, Fletcher B, Vandendriessche T, Ivics Z, Izsvak Z: Molecular evolution of a novel hyperactive Sleeping Beauty transposase

- enables robust stable gene transfer in vertebrates. *Nat Genet* 2009, **41**:753-761.
6. Liu G, Aronovich EL, Cui Z, Whitley CB, Hackett PB: **Excision of Sleeping Beauty transposons: parameters and applications to gene therapy.** *J Gene Med* 2004, **6**:574-583.
 7. Newman M, Wilson L, Camp E, Verdile G, Martins R, Lardelli M: **A Zebrafish Melanophore Model of Amyloidbeta Toxicity.** *Zebrafish* .
 8. Geurts AM, Yang Y, Clark KJ, Liu G, Cui Z, Dupuy AJ, Bell JB, Largaespada DA, Hackett PB: **Gene transfer into genomes of human cells by the sleeping beauty transposon system.** *Mol Ther* 2003, **8**:108-117.
 9. Yant SR, Park J, Huang Y, Mikkelsen JG, Kay MA: **Mutational analysis of the N-terminal DNA-binding domain of sleeping beauty transposase: critical residues for DNA binding and hyperactivity in mammalian cells.** *Mol Cell Biol* 2004, **24**:9239-9247.
 10. Zayed H, Izsvak Z, Walisko O, Ivics Z: **Development of hyperactive sleeping beauty transposon vectors by mutational analysis.** *Mol Ther* 2004, **9**:292-304.
 11. Baus J, Liu L, Heggestad AD, Sanz S, Fletcher BS: **Hyperactive transposase mutants of the Sleeping Beauty transposon.** *Mol Ther* 2005, **12**:1148-1156.
 12. Kirkitadze MD, Kowalska A: **Molecular mechanisms initiating amyloid beta-fibril formation in Alzheimer's disease.** *Acta Biochim Pol* 2005, **52**:417-423.
 13. Jowett T: **Tissue in situ hybridization.** John Wiley and Sons, New York; 1997.
 14. Higashijima S, Okamoto H, Ueno N, Hotta Y, Eguchi G: **High-frequency generation of transgenic zebrafish which reliably express GFP in whole muscles or the whole body by using promoters of zebrafish origin.** *Dev Biol* 1997, **192**:289-299.
 15. Kawakami K, Takeda H, Kawakami N, Kobayashi M, Matsuda N, Mishina M: **A transposon-mediated gene trap approach identifies developmentally regulated genes in zebrafish.** *Dev Cell* 2004, **7**:133-144.

doi:10.1186/1756-0500-3-282

Cite this article as: Newman and Lardelli: **A hyperactive sleeping beauty transposase enhances transgenesis in zebrafish embryos.** *BMC Research Notes* 2010 **3**:282.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

