

RESEARCH ARTICLE

Open Access

Methicillin-resistant staphylococcal contamination of cellular phones of personnel in a veterinary teaching hospital

Timothy Julian¹, Ameet Singh^{2*}, Joyce Rousseau¹ and J Scott Weese¹

Abstract

Background: Hospital-associated infections are an increasing cause of morbidity and mortality in veterinary patients. With the emergence of multi-drug resistant bacteria, these infections can be particularly difficult to eradicate. Sources of hospital-associated infections can include the patients' own flora, medical staff and inanimate hospital objects. Cellular phones are becoming an invaluable feature of communication within hospitals, and since they are frequently handled by healthcare personnel, there may be a potential for contamination with various pathogens. The objective of this study was to determine the prevalence of contamination of cellular phones (hospital issued and personal) carried by personnel at the Ontario Veterinary College Health Sciences Centre with methicillin-resistant *Staphylococcus pseudintermedius* (MRSP) and methicillin-resistant *Staphylococcus aureus* (MRSA).

Results: MRSP was isolated from 1.6% (2/123) and MRSA was isolated from 0.8% (1/123) of cellular phones. Only 21.9% (27/123) of participants in the study indicated that they routinely cleaned their cellular phone.

Conclusions: Cellular phones in a veterinary teaching hospital can harbour MRSP and MRSA, two opportunistic pathogens of significant concern. While the contamination rate was low, cellular phones could represent a potential source for infection of patients as well as infection of veterinary personnel and other people that might have contact with them. Regardless of the low incidence of contamination of cellular phones found in this study, a disinfection protocol for hospital-issued and personal cellular phones used in veterinary teaching hospitals should be in place to reduce the potential of cross-contamination.

Keywords: Cellular phone, Methicillin-resistant staphylococcus, Veterinary, Contamination

Background

Nosocomial or hospital-associated infections (HAIs) are an increasing cause of morbidity and mortality in human and veterinary medicine [1-4]. Multi-drug resistant (MDR) bacteria are commonly implicated in HAIs and can be challenging to eliminate [1-4]. Sources of HAI can include medical staff, the patients' own flora and inanimate hospital objects [5,6]. Hands of healthcare personnel are commonly contaminated with opportunistic pathogens and poor hand hygiene compliance is thought to be an important factor in the pathogenesis of HAIs [7]. Contaminated hands can result in direct transfer of pathogens to patients,

as well as contamination of inanimate objects (fomites). Any items that have frequent hand contact, especially in the absence of routine hand hygiene practices, are at high risk of becoming contaminated. Cellular phones (CPs) have become an indispensable accessory of today's society and they are being used extensively in a hospital setting to optimize patient care and client communications. However, CPs are commonly handled (irrespective of the cleanliness of hands), rarely disinfected and could harbour pathogenic bacteria. Concerns regarding bacterial contamination associated with the use of CPs within the hospital environment have been raised in human medicine, and studies of human healthcare worker CPs have reported contamination of 9–43% of CPs with bacteria known to cause HAIs [8-18]. Comparable data are not available for veterinary medicine.

* Correspondence: amsingh@uoguelph.ca

²Department of Clinical Studies, Ontario Veterinary College, University of Guelph, Guelph, ON N1G 2W1, Canada

Full list of author information is available at the end of the article

Methicillin-resistant *S. aureus* (MRSA) is a critically important hospital-associated pathogen in humans [19] and has been found on 1.9–10% of CPs sampled in hospitals [8-18]. It is also a significant concern in companion animals, both as a cause of HAI [1-3,20] and the potential for zoonotic transmission to veterinary personnel [21,22]. Of greater relevance from an animal health aspect is methicillin-resistant *S. pseudintermedius*, which has rapidly emerged as a leading cause of various opportunistic infections, including pyoderma [23,24] and surgical site infection [1-3].

The purpose of this study was to determine the MRSA and MRSP contamination rate of CPs used by personnel in the Ontario Veterinary College Health Sciences Centre (OVCHSC) and to identify factors associated with contamination. In this study we used electrostatic cloths to recover bacteria from CPs. This methodology has previously been described for the recovery of bacteria from inanimate objects [25-27]. This method was chosen because electrostatic cloths are easy to use, inexpensive and readily available. Furthermore, a standardized sampling technique for the recovery of bacteria from inanimate objects does not exist and the sensitivity of bacterial recovery from various techniques such as contact plates, electrostatic cloths and cotton-tipped swabs is unknown.

Methods

Sample collection and study population

This cross-sectional study was conducted from August to September 2011 at the OVCHSC. Hospital personnel, including veterinary students, technicians, residents/interns and clinical faculty were recruited for participation in this study. Upon verbal consent, participants' CPs (hospital-issued and/or personal) were sampled and a self-administered questionnaire was completed. The questionnaire characterized the participants' position in the hospital, frequency of CP cleaning, and whether the personal CP (if present) was used while in the OVCHSC. This study was approved by the University of Guelph Research Ethics Board.

Microbiological analysis

Samples were collected by wiping an electrostatic cloth (Swiffer[®], Procter & Gamble, Cincinnati, OH, USA) along all surfaces of the CP using a gloved hand [25-27]. After sampling, the cloth was placed into a pre-labeled sterile bag for later inoculation. All samples were collected by the same individual (TJ) and gloves were changed between each sample. To investigate potential sources of sampling contamination, electrostatic cloth samples were also collected from the hands and forearms of the individual (TJ) performing the sampling, the box containing the gloves, the clipboard and questionnaire sheets. During each sampling period, new, unused,

electrostatic cloths were removed from the box and cultured as negative controls [25].

Enrichment culture was performed by adding 70–80 mL of enrichment broth consisting of 10 g tryptone/L, 75 g sodium chloride/L, 10 g mannitol/L, and 2.5 g yeast extract/L to sterile bags containing electrostatic cloths immediately following sample collection. After 24 h incubation at 35°C, 1 loopful (approximately 10 µL) of broth was inoculated onto MRSA Chromogenic agar (BBL CHROMagar MRSA, Becton, Dickinson and Co., Mississauga, ON, Canada) and Mannitol Salt Agar with 2 µg/mL oxacillin and incubated at 35°C for MRSP selection. Plates were examined after 24 and 48 h of incubation.

Isolates were identified as *S. aureus* by colony morphology, pink color, Gram stain appearance, catalase reaction, coagulase reaction, and *S. aureus* latex agglutination test (Pastorex Staph-plus, Bio-Rad Laboratories Ltd, Mississauga, ON, Canada). Methicillin-resistance was confirmed by penicillin binding protein 2a latex agglutination test (MRSA latex agglutination test, Denka Seiken, USA Inc, Campbell, CA, USA). *Staphylococcus pseudintermedius* was identified by colony morphology, Gram stain appearance, catalase and coagulase reactions and a species-specific non-commercialized PCR assay [28].

MRSA characterization

Isolates were typed by sequencing of the X region of the protein A gene (*spa* typing) as described [29]. For *spa* typing, sequences were analyzed using eGenomics software (<http://tools.egenomics.com>). Ridom database equivalents were identified using the Ridom Spaserver website (<http://www.spaserver.ridom.de>). eGenomics *spa* types are reported using a numerical system (i.e. *spa* type 539) while Ridom *spa* types are reported using a numerical system preceded by a "t" (i.e. *spa* t034). Real-time PCR was used to detect the *lukF* and *lukS* components of the Pantone-Valentine leukocidin (PVL) using PVLSC-F (5'-GCTCAGGAGA TACAAG-3') and PVLSC-R (5'-GGATAGCAAAAG CAATG-3') primers [30].

MRSP characterization

MRSP isolates were characterized by sequence analysis of the *mec*-associated direct repeat unit (*dru* typing) [31], with *dru* repeats and types assigned by the Dru-typing.org database (<http://www.dru-typing.org/search.php>).

Statistical analysis

Contamination rate data were described. Categorical associations were assessed using Fisher's exact test (FET), with significance set at $P < 0.05$. Statistical analysis was performed using commercially available software (InStat, GraphPad Software Inc., La Jolla, CA, USA)

Results

One hundred and twenty three CPs from 106 individuals were sampled. Seventy-one were personal CPs while 52 were hospital-issued CPs (Table 1). Methicillin-resistant staphylococci were isolated from 3/123 (2.4%) CPs (Table 1). MRSP was isolated from two (1.6%) CPs while MRSA was isolated from one (0.8%). Both MRSP positive CPs were from samples collected on the same day and both were classified as dt9a. Samples were collected in an anonymous manner, with only identification of the personnel group (e.g. technician) so it was impossible to retrospectively investigate any potential associations between the two MRSP positive samples. Methicillin-resistant staphylococci were not isolated from any negative control samples.

The MRSA isolate was identified as *spa* type 18/t338 [32]. Based on available Canadian data [32] and Ridom Spaserver website (<http://www.spaserver.ridom.de>), this strain was inferred to be a clonal complex 30 strain that is classified as Canadian epidemic MRSA (CMRSA)-4, USA200, and eMRSA-16. It did not contain genes encoding PVL. The MRSP isolates were identified as *dru* type 9a, typically associated with sequence type 71 [33].

There was no difference in contamination between different personnel classifications ($P=0.2$, FET) overall, however there was a significant difference between groups when only personal CPs were considered ($P=0.004$, FET) but not with hospital-issued phones ($P=0.6$, FET). There was no difference overall between personal and hospital-issued phones ($P=1.0$, FET).

Twenty-two percent (27/123) of sampled CPs had reportedly been disinfected in the past month. Two/27 (7.4%) of the CPs that were contaminated had reportedly been disinfected in the past month versus 1/96 (1.0%) that had not been disinfected ($P=0.1$, FET). Sixty-four of the 71 (90%) of individuals carrying personal CPs reported using them while on the OVCHSC premises.

Table 1 MRSA and MRSP contamination rate of personal and hospital-issued CPs by veterinary hospital personnel

Group	Personal phones	Hospital-issued phones	Total
Technician	2/8 (25%) (1 MRSA, 1 MRSP)	0/15	2/23 (8.7%)
Residents/Interns	0/8	1/18 (5.6%) (MRSP)	1/23 (4.3%)
Veterinary student	0/37	0	0/37
Faculty	0/10	0/13	0/23
Non-medical personnel	0/8	0/6	0/14
Total	2/71 (2.8%)	1/52 (1.9%)	3/123 (2.4%)

Discussion

Contamination of CPs was uncommon but, nonetheless, both MRSA and MRSP were identified. These results support concerns that CPs could act as a fomite for pathogenic bacteria, with transmission to patients or personnel through subsequent contamination of the hands. The relevance of CP contamination is unclear but these data, along with similar data from human medicine raise concern [8-18]. People frequently handle CPs, and likely do so irrespective of the cleanliness of their hands. Goldblatt *et al.* reported that physicians used CPs excessively, even during patient contact, which likely contributed to a higher rate of CP contamination compared with nursing staff [18]. These factors create the potential for both contamination of the phone from contaminated hands, and transfer of pathogens from a contaminated CP to clean hands. Proper hand hygiene has been emphasized as a means of reducing the incidence of nosocomial infections [7] and is probably a key factor for reducing CP contamination. If hand hygiene is properly performed before and after patient contact, and after contact with potentially contaminated environmental sites, the risk of CP contamination would be minimized. However, hand hygiene compliance rates in human medicine remains below 50% [34] and there is no evidence suggesting better compliance in veterinary health-care personnel.

In a human healthcare study, physician CPs were more often contaminated than those of nursing staff (60% vs 20%) [18]. Here, there was no significant difference overall, however there was a significant difference between groups when only personal CPs were considered, with contamination identified only on technician CPs. Reasons for this are unclear and were not specifically investigated here. It is possible that contamination could be of greater risk in technicians because they may have more contact with animals and handle more animals overall compared to the other groups. At the OVCHSC, personnel use their personal CP in the hospital and community, whereas hospital-issued CPs are used only within hospital premises. Although there was no significant difference in contamination between personal and hospital-issued CPs, further study of this association and other factors associated with CP contamination are required.

MRSP is a significant problem in companion animals, and the presence of this multidrug opportunistic pathogen is a concern [35-38]. This is not surprising since MRSP can be found in clinically normal animals [39] and in the veterinary hospital environment [40]. Considering the significant problems encountered in treating some MRSP infections, especially implant-associated surgical site infections, measures to reduce transmission of this pathogen are important. The MRSP isolates identified here were dt9a, which is among the most common

dru types in North America and typically associated with sequence type 71 [33].

MRSA is both an animal health and zoonotic concern. The isolate identified here was *spa* type 18/t338, a human epidemic clone. Typing cannot indicate the origin of MRSA because companion animals are typically infected with human MRSA clones, and no testing of personnel was performed to differentiate animal versus human sources [20].

Only 21.9% of sampled CPs had been disinfected within the past month. Method of disinfection was not queried in the survey administered to participants. Facility infection control protocols do not address disinfection of CPs and that, combined with lack of consideration of the potential for CPs to become contaminated, may explain the low disinfection rates. There is also limited information about CP disinfection methods that are both effective and do not damage the CP. However, the use of 70% isopropyl alcohol wipes eliminated bacterial contamination in 98% of mobile phones in one study [41], and this is a simple measure that could be used routinely.

Conclusions

While uncommon, contamination of CPs with MRSA and MRSP was identified, supporting concerns that these devices may be fomites for transmission of infection to patients or personnel. Further, since personal CPs are used in the hospital and community, they represent a potential bridge between the hospital and community and could transfer zoonotic pathogens to anyone that has contact with the CP. The true risk of CP contamination is not known and whether these fomites play a role in transmission of MRSA and/or MRSP requires further investigation. Increased consideration should be given to reducing contamination, particularly avoiding handling CPs when hands might be contaminated and using good hand hygiene practices. Routine disinfection of CPs, such as with alcohol wipes, while unproven in a veterinary context, should be considered as part of a general infection control program.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Pathobiology, Ontario Veterinary College, University of Guelph, Guelph, ON N1G 2W1, Canada. ²Department of Clinical Studies, Ontario Veterinary College, University of Guelph, Guelph, ON N1G 2W1, Canada.

Authors' contributions

TJ performed data collection and drafted the manuscript. TJ and JR performed microbiological testing. AS and JSW were responsible for study design and manuscript review. All authors read and approved the final manuscript.

Received: 07 December 2011 Accepted: 25 April 2012

Published: 25 April 2012

References

1. Faires MC, Traverse M, Tater KC, Pearl DL, Weese JS: **Methicillin-resistant and -susceptible *Staphylococcus aureus* infections in dogs.** *Emerg Infect Dis* 2010, **16**:69–75.
2. Weese JS: **A review of multidrug resistant surgical site infections.** *Vet Comp Orthop Traumatol* 2008, **21**:1–7.
3. Weese JS: **A review of post-operative infections in veterinary orthopedic surgery.** *Vet Comp Orthop Traumatol* 2008, **21**:99–105.
4. Owens CD, Stoessel K: **Surgical site infections: epidemiology, microbiology and prevention.** *J Hosp Infect* 2008, **70**(Suppl 2):3–10.
5. Hardy KJ, Oppenheim BA, Gossain S, Gao F, Hawkey PM: **A study of the relationship between environmental contamination with methicillin-resistant *Staphylococcus aureus* (MRSA) and patients' acquisition of MRSA.** *Infect Control Hosp Epidemiol* 2006, **27**:127–132.
6. Marshal BM, Ochieng DJ, Levy SB: **Commensals: unappreciated reservoir of antibiotic resistance.** *Microbe* 2009, **4**:231–238.
7. Allegranzi B, Pittet D: **Role of hand hygiene in healthcare associated infection prevention.** *J Hosp Infect* 2009, **73**:305–315.
8. Brady RRW, Verran J, Daman NN, Gibb AP: **Review of mobile communication devices as potential reservoirs of nosocomial pathogens.** *J Hosp Inf* 2009, **41**:295–300.
9. Brady RRW, Fraser SF, Dunlop MG, Paterson-Brown S, Gibb AP: **Bacterial contamination of mobile communication devices in the operative environment.** *J Hosp Infect* 2007, **66**:397–398.
10. Karabay O, Kocoglu E, Tahtaci M: **The role of mobile phones in the spread of bacteria associated with nosocomial infections.** *J Infect Developing Countries* 2007, **1**:72–73.
11. Khivsara A, Sushma T, Dhanashree B: **Typing of *Staphylococcus aureus* from mobile phones and clinical samples.** *Curr Sci* 2006, **90**:910–912.
12. Jeske HC, Tiefenthaler W, Hohlrieder M, Hinterberger G, Benzer A: **Bacterial contamination of anaesthetist's hands by personal mobile phone and fixed phone use in the operating theatre.** *Anaesthesia* 2007, **62**:904–906.
13. Borer A, Gilad J, Smolyakov R: **Cell phones and Acinetobacter transmission.** *Emerg Infect Dis* 2005, **11**:1160–1161.
14. Jayalakshmi J, Appalaraju B, Usha S: **Cell phones as reservoir of nosocomial pathogens.** *J Assoc Physicians India* 2008, **56**:388–389.
15. Hassoun A, Vellozi EM, Smith MA: **Colonization of personal digital assistants carried by healthcare professionals.** *Infect Control Hosp Epidemiol* 2004, **25**:1000–1001.
16. Tekerekoglu MS, Duman Y, Serindag A, Cuglan SS, Kaysadu H, Tunc E, Yakupogullari Y: **Do mobile phones of patients, companions and visitors carry multidrug-resistant hospital pathogens?** *Am J Infect Control* 2011, **39**:379–381.
17. Sadat-Ali M, Al-Omran AK, Azam Q, Bukari H, Al-Zahrani AJ, Al-Turki RA, Al-Omran AS: **Bacterial flora on cell phones of health care providers in a teaching institution.** *Am J Infect Control* 2010, **38**:404–405.
18. Goldblatt JG, Krief I, Klonsky T: **Use of cellular telephones and transmission of pathogens by medical staff in New York and Israel.** *Infect Control Hosp Epidemiol* 2007, **28**:500–503.
19. Klevens RM, Morrison MA, Nadle J, Petit S, Gershman K, Ray S, Harrison LH, Lynfield R, Dumyati G, Townes JM, Craig AS, Zell ER, Fosheim GE, McDougal LK, Carey RB, Fridkin SK, Active Bacterial Core surveillance MRSA investigators: **Invasive methicillin-resistant *Staphylococcus aureus* infections in the United States.** *JAMA* 2007, **298**:1763–1771.
20. Weese JS, van Duijkeren E: **Methicillin-resistant *Staphylococcus aureus* and *Staphylococcus pseudintermedius* in veterinary medicine.** *Vet Microbiol* 2010, **140**:418–429.
21. Loeffler A, Pfeiffer DU, Lloyd DH, Smith H, Soares-Magalhaes R, Lindsay JA: **Methicillin-resistant *Staphylococcus aureus* carriage in UK veterinary staff and owners of infected pets: new risk groups.** *J Hosp Infect* 2010, **74**:282–288.
22. Burstiner LC, Faires M, Weese JS: **Methicillin-resistant *Staphylococcus aureus* colonization in personnel attending a veterinary surgery conference.** *Vet Surg* 2010, **39**:150–157.
23. Griffith GC, Morris DO, Abraham JL, Shofer FS, Rankin SC: **Screening for skin carriage of methicillin-resistant coagulase-positive staphylococci and *Staphylococcus schleiferi* in dogs with healthy and inflamed skin.** *Vet Dermatol* 2008, **19**:142–149.

24. Onuma K, Tanabe T, Sato H: **Antimicrobial resistance of *Staphylococcus pseudintermedius* isolates from healthy dogs and dogs affected with pyoderma in Japan.** *Vet Dermatol* 2011. doi:10.1111/j.1365-3164.2011.00995.x [Epub ahead of print].
25. Burgess BA, Morley PS, Hyatt DR: **Environmental surveillance for *Salmonella enterica* in a veterinary teaching hospital.** *J Am Vet Med Assoc* 2004, **225**:1344–1348.
26. Hoet AE, Johnson A, Nava-Hoet RC, Bateman S, Hillier A, Dyce J, Gebreyes WA, Wittum TE: **Environmental methicillin-resistant *Staphylococcus aureus* in a veterinary teaching hospital during a nonoutbreak period.** *Vect Zoonot Dis* 2011, **11**:609–615.
27. Murphy CP, Reid-Smith RJ, Boerlin P, Weese JS, Prescott JF, Janecko N, Hassard L, McEwen SA: ***Escherichia coli* and selected veterinary and zoonotic pathogens isolated from environmental sites in companion animal veterinary hospitals in southern Ontario.** *Can Vet J* 2010, **51**:963–972.
28. Sasaki T, Tsubakishita S, Tanaka Y, Sakusabe A, Ohtsuka M, Kawakami T, Fukata T, Hiramatsu K: **Multiplex-PCR method for species identification of coagulase-positive staphylococci.** *J Clin Microbiol* 2010, **48**:765–769.
29. Shopsis B, Gomez M, Montgomery SO, Smith DH, Waddington M, Dodge DE, Bost DA, Riehman M, Naidich S, Kreiswirth BN: **Evaluation of protein A gene polymorphic region DNA sequencing for typing of *Staphylococcus aureus* strains.** *J Clin Microbiol* 1999, **37**:3556–3563.
30. Rankin S, Roberts S, O'Shea K, Maloney D, Lorenzo M, Benson CE: **Panton-Valentine leukocidin (PVL) toxin positive MRSA strains isolated from companion animals.** *Vet Microbiol* 2005, **108**:145–148.
31. Goering RV, Morrison D, Al-Doori Z, Edwards GF, Gemmell CG: **Usefulness of mec-associated direct repeat unit(dru) typing in the epidemiological analysis of highly clonal methicillin-resistant *Staphylococcus aureus* in Scotland.** *Clin Microb Infect* 2008, **14**:964–969.
32. Golding GR, Campbell JL, Spreitzer DJ, Veyhl J, Surynicz K, Simor A, Mulvey MR, Canadian Nosocomial Infection Surveillance Program: **A preliminary guideline for the assignment of methicillin-resistant *Staphylococcus aureus* to a Canadian pulsed-field electrophoresis epidemic type using spa typing.** *Can J Infect Dis Med Microbiol* 2008, **9**:273–281.
33. Kadlec K, Schwarz S, Weese JS: **Evaluation of direct repeat unit (dru) typing for characterization of methicillin-resistant *Staphylococcus pseudintermedius*.** In *2nd ASM/ESCMID conference on methicillin-resistant staphylococci in animals: veterinary and public health implications: 8–11 September 2011; Washington*
34. McGuckin M, Waterman R, Govednik J: **Hand hygiene compliance rates in the United States – a one-year multicenter collaboration using product/volume usage measurement and feedback.** *Am J Med Qual* 2009, **24**:205–213.
35. van Duijkeren E, Houwers DJ, Schoormans A, Broekhuizen-Stins MJ, Ikawaty R, Fluit AC, Wagenaar JA: **Transmission of methicillin-resistant *Staphylococcus intermedius* between humans and animals.** *Vet Microbiol* 2008, **128**:213–215.
36. Laarhoven LM, de Heus P, van Luijn J, Duim B, Wagenaar JA, van Duijkeren E: **Longitudinal study on methicillin-resistant staphylococcus pseudintermedius in Households.** *PLoS One* 2011, **6**:e27788.
37. Frank LA, Kania SA, Kirzeder EM, Eberlein LC, Bemis DA: **Risk of colonization or gene transfer to owners of dogs with methicillin-resistant *Staphylococcus pseudintermedius*.** *Vet Dermatol* 2009, **20**:496–501.
38. Perreten V, Kadlec K, Schwarz S, Grönlund Andersson U, Finn M, Greko C, Moodley A, Kania SA, Frank LA, Bemis DA, Franco A, Iurescia M, Battisti A, Duim B, Wagenaar JA, van Duijkeren E, Weese JS, Fitzgerald JR, Rossano A, Guardabassi L: **Clonal spread of methicillin-resistant *Staphylococcus pseudintermedius* in Europe and North America: an international multi-center study.** *J Antimicrob Chemother* 2010, **65**:1145–1154.
39. Murphy C, Reid-Smith RJ, Prescott JF, Bonnett BN, Poppe C, Boerlin P, Weese JS, Janecko N, McEwen SA: **Occurrence of antimicrobial resistant bacteria in healthy dogs and cats presented to private veterinary hospitals in southern Ontario: a preliminary study.** *Can Vet J* 2009, **50**:1047–1053.
40. Nienhoff U, Kadlec K, Chaberny IF, Verspohl J, Gerlach GF, Kreienbrock L, Schwarz S: **Methicillin-resistant *Staphylococcus pseudintermedius* among dogs admitted to a small animal hospital.** *Vet Microbiol* 2011, **150**:191–197.
41. Arora U, Devi P, Chadha A, Malhotra S: **Cellphones a modern stayhouse for bacterial pathogens.** *JK Science* 2009, **11**:127–129.

doi:10.1186/1756-0500-5-193

Cite this article as: Julian et al.: Methicillin-resistant staphylococcal contamination of cellular phones of personnel in a veterinary teaching hospital. *BMC Research Notes* 2012 **5**:193.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

