

RESEARCH ARTICLE

Open Access

# Antibacterial and antibiotic-potential activities of the methanol extract of some cameroonian spices against Gram-negative multi-drug resistant phenotypes

Igor K Voukeng<sup>1</sup>, Victor Kuete<sup>1\*</sup>, Jean P Dzoyem<sup>1</sup>, Aimé G Fankam<sup>1</sup>, Jaures A K Noumedem<sup>1</sup>, Jules R Kuate<sup>1</sup> and Jean-Marie Pages<sup>2</sup>

## Abstract

**Background:** The present work was designed to evaluate the antibacterial properties of the methanol extracts of eleven selected Cameroonian spices on multi-drug resistant bacteria (MDR), and their ability to potentiate the effect of some common antibiotics used in therapy.

**Results:** The extract of *Cinnamomum zeylanicum* against *Escherichia coli* ATCC 8739 and AG100 strains showed the best activities, with the lowest minimal inhibitory concentration (MIC) of 64 µg/ml. The extract of *Dorstenia psilurus* was the most active when tested in the presence of an efflux pump inhibitor, phenylalanine Arginine-β-Naphtylamide (PAβN), a synergistic effect being observed in 56.25 % of the tested bacteria when it was combined with Erythromycin (ERY).

**Conclusion:** The present work evidently provides information on the role of some Cameroonian spices in the fight against multi-resistant bacteria.

**Keywords:** Multi-Drug Resistant bacteria, Spices, Methanol extract, Cameroon

## Background

Infectious diseases are one of the leading causes of morbidity and mortality worldwide, especially in developing countries [1,3]. Following the massive use of antibiotics in human therapy, bacteria have developed several resistance mechanisms including the efflux of antibiotics [3]. Several Cameroonian spices are known to possess medicinal values [4]. In our previous report, we demonstrated that several medicinal spices inhibited the growth of MDR bacteria and were also able to improve the activity of commonly used antibiotics [5]. In our continuous search of antimicrobial drugs from medicinal plant, we designed the present work to investigate the antibacterial potential against Gram-negative MDR bacteria of some of the commonly used medicinal spices in Cameroon

such as *Aframomum citratum* (Pereira) K. Schum. (Zingiberaceae), *Aframomum melegueta* (Roscoe) K. Schum. (Zingiberaceae), *Scorodophloeus zenkeri* Harms (Caesalpinaceae), *Tetrapleura tetraptera* (Schum. & Thonn) Taub. (Mimosaceae), *Fagara leprieurii* (Guill and Perr) Engl. (Rutaceae), *Monodora myristica* Dunal (Annonaceae), *Piper guineense* (Schum and Thonn) (Piperaceae), *Dorstenia psilurus* Welwitsch (Moraceae), *Imperata cylindrica* Beauv. var. *koenigii* Durand and Schinz (Gramineae), *Pentadiplandra brazzeana* Baill. (Capparaceae) and *Cinnamomum zeylanicum* (Linn) Cor. (Lauraceae).

## Material and methods

### Plant materials and extraction

The eleven edible spices used in this work were purchased from Dschang local market, West Region of Cameroon in January 2010. The collected spices material were the fruits of *Aframomum citratum*, *Aframomum melegueta*, *Scorodophloeus zenkeri*, *Tetrapleura tetraptera*,

\* Correspondence: kuetevictor@yahoo.fr

<sup>1</sup>Department of Biochemistry, Faculty of science, University of Dschang, P.O. Box 67, Dschang, Cameroon, Africa

Full list of author information is available at the end of the article

**Table 1 Spices used in the present study and evidence of their activities**

Spice samples (Family)	Herbarium Voucher number <sup>a</sup>	Traditional Treatment	Part used	Bioactive (or potentially active) compounds <sup>b</sup> and screened activity <sup>c</sup> for crude plant extract
<i>Aframomum citratum</i> (Pereira) K. Schum. (Zingiberaceae)	37 736/HNC	Malaria, aphrodisiac, cancer [4,6]	Fruits, leaves, seeds	<u>Antimicrobial</u> : Ethylacetate extract of fruits on <i>Ec. Pa. Sa</i> [7]  <u>Cytotoxicity of fruits crude methanol extract</u> [weak activity on leukemia CCRF-CEM and CEM/ADR5000 cells, and pancreatic MiaPaCa-2 cell lines] [4]
<i>Aframomum melegueta</i> (Roscoe) K. Schum. (Zingiberaceae)	39 065/HNC	Malaria, dysentery, carminative, dysmenorrheal, fertility, rubella, leprosy, cancer [6,8]	Fruits, leaves	<u>Antimicrobial</u> : Aqueous and ethanol extract of leaves on <i>Fo. An</i> [9] Methanol extract of fruits ( <i>Q</i> ) on <i>Sa. Bs. Ec. Pa. Ca</i> [8]  <u>Cytotoxicity of fruits crude methanol extract</u> [weak activity on leukemia CCRF-CEM and pancreatic MiaPaCa-2 cell lines and significant activity on CEM/ADR5000 cells with IC <sub>50</sub> value of 7.08 µg/ml] [4]
<i>Cinnamomum zeylanicum</i> (Linn) Cor. (Lauraceae)	22 309/SRFC	Cancer [4]	Fruits, leaves, bark	<u>Antimicrobial</u> : <i>Cd, Cm, Lt, Fp</i> [10,11]  <u>Cytotoxicity of leaves crude methanol extract</u> [weak activity on leukemia CCRF-CEM and CEM/ADR5000 cells, and pancreatic MiaPaCa-2 cell lines] [4]
<i>Dorstenia psilurus</i> Welwitsch (Moraceae)	44 839/HNC	Snake bite, rheumatism, head and stomach ache, hypertension, cancer [4,12,13].	Leaves, roots	<u>Cytotoxicity of roots crude methanol extract</u> [Significant activity with IC <sub>50</sub> values of 7.18; 7.79 and 9.17 µg/ml respectively on leukemia CCRF-CEM cells, CEM/5000 cells and pancreatic MiaPaCa-2 cell lines] [4]
<i>Fagara lepreurii</i> (Guill and Perr) Engl. (Rutaceae)	37 632/HNC	Gastritis, gingivitis, bilharzias, anti-diarrhoeal, cancer, laxative, antimicrobial, ulcer, gonorrhoea, kidney ache, sterility [4,14,15]	Bark, leaves, roots	<u>Antimicrobial</u> : Ethanol extract of the seeds on <i>Ca. Cn. Mg. Tm. Tr. Bci. Af. Af. Sb</i> [6] <u>Essential oil</u> : <i>Sa</i> [15]  <u>Cytotoxicity of seeds crude methanol extract</u> [weak activity on leukemia CCRF-CEM and pancreatic MiaPaCa-2 cell lines and significant activity on CEM/ADR5000 cells with IC <sub>50</sub> value of 8.13 µg/ml] [4]
<i>Imperata cylindrica</i> Beauv. var. <i>koenigii</i> Durand et Schinz (ramineae)	30 139/SRFC	Diuretic, anti-inflammatory, dysentery, urinary tract infections, cancer [4,16,17]	Leaves, roots	<u>Cytotoxicity of roots crude methanol extract</u> [Significant activity with IC <sub>50</sub> values of 8.4; 7.18 and 12.11 µg/ml respectively on leukemia CCRF-CEM cells, CEM/5000 cells and pancreatic MiaPaCa-2 cell lines] [4]
<i>Monodora myristica</i> Dunal (Annonaceae)	2 949/SRFC	Insecticidal, diuretic, constipation, anti-hemorrhage, headache, wounds, worm infections, cancer [4,15,18,19]	Fruits, leaves, seeds	<u>Antimicrobial</u> : <i>Fm. Af. Af</i> [18]; <u>Essential oil</u> : <i>Af. Bc. Bs. Cgl. Ec. Kp. Sa. Sf</i> [15]. <u>Cytotoxicity of fruits seeds methanol extract</u> [weak activity on leukemia CCRF-CEM and CEM/ADR5000 cells, and pancreatic MiaPaCa-2 cell lines] [4]
<i>Pentadiplandra brazeana</i> Baill. (Capparaceae)	42 918/HNC	Gastric ulcer, cancer [4,20]	Fruits, leaves	<u>Cytotoxicity of roots crude methanol extract</u> [weak activity on leukemia CCRF-CEM and CEM/ADR5000 cells, and pancreatic MiaPaCa-2 cell lines] [4]
<i>Piper guineense</i> (Schum and Thonn) (Piperaceae)	6 018/SRFC	Cough, bronchitis, rheumatism, insecticidal, anemia, carminative, stomach ache, cancer [4,8,21]	Fruits, leaves, bark	<u>Insecticidal</u> : <i>Cs</i> [20] <u>Antimicrobial</u> ( <i>Q</i> ); <i>Ec. Sa. Bs. Pa. Ca. An</i> [8,22]
<i>Scorodophloeus zenkerii</i> Harms (Caesalpinaceae)	44 803/HNC	Cancer [4]	Leaves, roots	<u>Antimicrobial</u> : Essential oil of stem bark on <i>Ec, Sa, Bs, Cu</i> [23].  <u>Cytotoxicity of fruits crude methanol extract</u> [weak activity on leukemia CCRF-CEM and CEM/ADR5000 cells, and pancreatic MiaPaCa-2 cell lines] [4]

**Table 1 Spices used in the present study and evidence of their activities (Continued)**

<i>Tetrapleura tetraptera</i> (Schum. & Thonn) Taub. (Mimosaceae)	12 117/SRFC	Pain, arthritis, epilepsy, convulsion, gastric ulcer, cancer [4,20]	Bark, leaves, roots	Cytotoxicity of fruits crude methanol extract [weak activity on leukemia CCRF-CEM and CEM/ADR5000 cells, and pancreatic MiaPaCa-2 cell lines] [4]
---	-------------	---	---------------------	---

<sup>a</sup>(HNC): Cameroon National Herbarium; (SRFC): *Société des réserves forestières du Cameroun*; <sup>b</sup>(/): Not reported.

<sup>c</sup>[Screened activity: significant (S: CMI < 100 µg/ml). moderate (M : 100 < CMI ≤ 625 µg/ml). Weak (W: CMI > 625 µg/ml) Q: Qualitative activity based on the determination of inhibition zone [15]; Af : *Aspergillus fumigatus*. Afl : *Aspergillus flavus*. An : *Aspergillus niger*. Bc : *Bacillus cereus*. Bci : *Botrytis cinerea*. Bs : *Bacillus subtilis*. Bt : *Botrydiploidea theobromae*. Ca : *Candida albicans*. Cd : *Clostridium difficile*. Cm : *Colletotrichum musae*. Cn : *Cryptococcus neoformans*. Cs : *Callosobruchus subinnotatus*. Cu : *Candida utilis*. Ec : *Escherichia coli*. Fm : *Fusarium moniliforme*. Fo : *Fusarium oxysporum*. Fp : *Fusarium proliferatum*. Lt : *Lasiodiplodia theobromae*. Mg : *Microsporium gypseum*. Pa: *Pseudomonas aeruginosa*. Sa : *Staphylococcus aureus*. Sb: *Scopulariopsis brevicaulis*. Tm: *Trichophyton mentagrophytes*. Tr: *Trichophyton rubrum*.

the seeds of *Fagara leuprieurii*, *Monodora myristica* and *Piper guineense*, the roots of *Dorstenia psilurus*, *Imperata cylindricum* and *Pentadiplandra brazzeana* and the leaves of *Cinnamomum zeylanicum*. The plants were identified by Mr. Victor Nana of the National herbarium (Yaoundé, Cameroon) where voucher specimens were deposited under a reference number (Table 1). The extracts were obtained by methanol (MeOH) maceration as previously described [5].

#### Preliminary phytochemical investigations

The major secondary metabolites classes were screened according to the common phytochemical methods described by Harborne [24].

#### Chemicals for antimicrobial assays

Tetracycline (TET), cefepime (FEP), streptomycin (STR), ciprofloxacin (CIP), norfloxacin (NOR), chloramphenicol (CHL), cloxacillin (CLX), ampicillin (AMP), erythromycin (ERY), kanamycin (KAN) (Sigma-Aldrich, St Quentin Fallavier, France) were used as reference

antibiotic. *p*-Iodonitrotetrazolium chloride (INT) and phenylalanine arginine β-naphthylamide (PAβN) were used as microbial growth indicator and efflux pumps inhibitor (EPI) respectively.

#### Bacterial strains and culture media

The studied microorganisms included reference (from the American Type Culture Collection) and clinical (Laboratory collection) strains of *Providencia stuartii*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Escherichia coli*, *Enterobacter aerogenes* and *Enterobacter cloacae*. The bacterial strains and their features were previously reported [5]. The preliminary treatment of these organisms as well as the culture media were conducted as previously described [5].

#### Bacterial susceptibility determinations

The respective MICs of samples on the studied bacteria were determined using rapid INT colorimetric assay [25,26] with some modifications as previously reported [5]. The inoculum concentration used was 1.5 x10<sup>6</sup> CFU/ml and the samples were incubated at 37 °C for

**Table 2 Extraction yields, aspects and phytochemical composition of the plant extracts**

Spice samples	Extraction yield (%) <sup>*</sup>	Physical aspect	Phytochemical composition								
			Alkaloids	Anthocyanins	Anthraquinones	Flavonoids	Phenols	Saponins	Sterols	Tannins	Triterpenes
<i>Aframomum citratum</i>	2.6	Oily, dark green	+	-	-	+	+	-	-	+	+
<i>Aframomum megueta</i>	7.3	Brown powder	+	-	-	-	-	+	-	-	+
<i>Cinnamomum zeylanicum</i>	8.4	Oily, dark green	+	-	-	+	+	-	+	+	-
<i>Dorstenia psilurus</i>	10.3	Oily, brown	+	+	+	+	+	+	-	+	+
<i>Fagara leuprieurii</i>	26.2	Creamy, brown	+	-	+	+	+	-	-	+	+
<i>Imperata cylindricum</i>	8.2	Creamy, brown	+	+	+	+	+	-	-	-	+
<i>Monodora myristica</i>	23.5	Oily, brown	+	-	+	+	+	-	-	-	+
<i>Pentadiplandra brazzeana</i>	4.6	Creamy, brown	+	-	-	+	+	-	-	-	-
<i>Piper guineense</i>	17.5	Creamy, brown	+	-	+	+	+	-	-	-	-
<i>Scorodophloeus zenkeri</i>	9.2	Creamy, dark green	+	-	-	+	+	-	-	+	-
<i>Tetrapleura tetraptera</i>	29.4	brown	+	-	+	+	+	+	-	+	+

(+): Present; (-): Absent; <sup>\*</sup>The yield was calculated as the ratio of the obtained methanol extract according to the initial mass of the spice powder.

**Table 3 Minimal inhibitory concentration (MIC) of the studied plants extracts and chloramphenicol on the studied bacterial species**

Bacterial strains	Tested samples and MIC in µg/ml in the absence and presence of PAβN (in parenthesis)											
	<i>Aframomum citratum</i>	<i>Aframomum melegueta</i>	<i>Imperata cylindricum</i>	<i>Cinnamomum zeylanicum</i>	<i>Dorstenia psilurus</i>	<i>Fagara leprieuri</i>	<i>Monodora myristica</i>	<i>Pentadiplandra brazzeana</i>	<i>Piper guineense</i>	<i>Scorodophloeus zenkeri</i>	<i>Tetrapleura tetraptera</i>	CHL
<b><i>E. coli</i></b>												
ATCC8739	512	512	512	<b>64</b>	-	512	1024	1024	1024	1024	1024	1
ATCC10536	1024	512	1024	512	128	256	1024	512	1024	512	1024 (1024)	32 (<2)
AG100	1024 (1024)	1024 (1024)	1024 (256)	- ( <b>64</b> )	- (1024)	1024 (1024)	512	1024 (1024)	1024 (512)	1024 (1024)	1024 (1024)	4 (<2)
AG100A	512 (128)	1024 (1024)	1024 (128)	512 (128)	512 (128)	512 (512)	1024 (1024)	- (-)	1024 (1024)	1024 (1024)	1024 (1024)	<2 (<2)
AG100A <sub>TET</sub>	512 (512)	1024 (1024)	1024 (1024)	512 (512)	512 (128)	1024 (1024)	-	-	1024	512	1024	32 (<2)
AG102	1024	-	1024	1024	512	1024	-	-	-	-	-	16 (<2)
MC4100	512 (512)	512 (256)	1024 (1024)	1024 (1024)	512 (256)	512	- (-)	1024 (1024)	1024 (1024)	1024 (1024)	512 (512)	4 (<2)
W3110	512 (256)	512 (512)	512 (512)	512 (512)	512 (256)	256	512	1024 (1024)	1024 (128)	512	512 (512)	1 (<2)
<b><i>E. aerogenes</i></b>												
ATCC13048	1024	-	1024	1024	1024	1024	1024	-	-	-	-	8 (<2)
CM64	1024 (1024)	1024 (1024)	512 (128)	1024 (512)	512 (256)	1024 (1024)	1024 (1024)	1024 (1024)	1024 (1024)	1024 (1024)	512 (512)	32
EA27	512 (512)	1024 (1024)	512 (512)	512 (512)	- (-)	1024 (1024)	1024 (1024)	1024 (1024)	- (-)	1024 (1024)	1024 (512)	64 (32)
EA289	-	1024	-	1024	-	-	1024	1024	-	-	1024	256
EA298	1024	512	-	-	1024	-	256	256	512	256	1024	256
EA3	-	-	-	-	-	1024	-	-	-	-	-	256
<b><i>E. cloacae</i></b>												
BM47	512 (512)	1024 (1024)	1024 (1024)	1024 (512)	1024 (128)	1024 (1024)	1024 (1024)	1024	1024	1024	1024	- (8)
BM67	512 (512)	1024 (1024)	1024 (1024)	1024 (1024)	1024 (128)	- (-)	- (-)	- (-)	- (-)	- (-)	- (-)	- (32)
ECCL69	512 (512)	1024 (1024)	1024 (1024)	1024 (1024)	- (-)	- (-)	1024 (1024)	- (-)	1024 (1024)	1024 (1024)	1024 (512)	- (32)
<b><i>K. pneumoniae</i></b>												
ATCC12296	1024	1024	1024	1024	1024	1024	512	-	-	-	1024	4
K2	1024	-	1024	1024	1024	-	1024	-	-	-	-	-
K24	1024	1024	1024	1024	1024	1024	512	-	-	1024	1024	32 (<2)

**Table 3 Minimal inhibitory concentration (MIC) of the studied plants extracts and chloramphenicol on the studied bacterial species (Continued)**

KP55	512	1024	256	512	512	1024	1024	-	-	-	1024	32 (<2)
KP63	512 (512)	1024 (1024)	1024 (1024)	512 (512)	512 (128)	1024 (512)	1024 (1024)	512	1024 (1024)	1024 (512)	1024 (1024)	64 (<2)
<b><i>P. stuartii</i></b>												
ATCC29916	1024 (1024)	- (-)	- (-)	- (-)	1024 (1024)	- (-)	- (-)	- (-)	1024 (1024)	1024 (1024)	1024 (1024)	8
NEA16	1024 (512)	- (-)	1024 (1024)	512 (512)	512 (256)	1024	1024	-	-	1024	-	64 (<2)
PS2636	1024	-	-	-	-	-	-	-	-	-	-	-
PS299645	512	512	1024	1024	1024	1024	-	1024	1024	512	1024	128
<b><i>P. aeruginosa</i></b>												
PA01	-	-	-	-	-	-	-	-	-	-	-	-
PA124	-	-	-	-	-	-	-	-	-	1024	-	32 (<2)

(-): MIC not detected at up to 1024 µg/ml for the les extracts and 256 µg/ml for chloramphenicol. ( ) : values in parenthesis are MIC of substance in the presence of PAβN at 20 µg/ml. The MIC of PAβN was 64 µg/ml on *E. coli*. AG100A. 512 µg/ml on ATCC11296. BM67. EA27. EA289; 1024 µg/ml on AG100A<sub>TET</sub>. ATCC13048. CM64; and > 1024 µg/ml on other bacteria. CHL: chloramphénicol; (in bold): significant MIC value.

**Table 4 Minimal inhibitory concentration (MIC) in µg/ml of antibiotics in the absence and presence sub-inhibitory concentrations of *Aframomum citratum* extract against some MDR bacteria**

Bacterial strains	Antibiotics and MIC in absence and presence of <i>Aframomum citratum</i> extract														
	Ampicillin			Cefepime			Chloramphenicol			Ciprofloxacin			Cloxacillin		
	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5
AG100Atet	-	-	-	-	-	-	256	32 (8) <sup>S</sup>	64 (4) <sup>S</sup>	256	128 (2) <sup>S</sup>	256 (1) <sup>I</sup>	-	-	-
AG102	-	-	-	128	128 (1) <sup>I</sup>	128 (1) <sup>I</sup>	16	8 (2) <sup>S</sup>	8 (2) <sup>S</sup>	<2	<2	<2	-	-	-
CM64	-	256 (1) <sup>I</sup>	-	-	64 (>4) <sup>S</sup>	-	nt	nt	nt	nt	nt	nt	-	-	-
KP63	-	32 (>8) <sup>S</sup>	-	256	32 (8) <sup>S</sup>	256 (1) <sup>I</sup>	-	64 (>4) <sup>S</sup>	256 (>1) <sup>S</sup>	64	64 (1) <sup>I</sup>	64 (1) <sup>I</sup>	-	256 (>1) <sup>S</sup>	-
PA124	128	16 (8) <sup>S</sup>	64 (2) <sup>S</sup>	128	128 (1) <sup>I</sup>	256 (0.5) <sup>I</sup>	32	16 (2) <sup>S</sup>	16 (2) <sup>S</sup>	16	4 (4) <sup>S</sup>	16 (1) <sup>I</sup>	-	-	-
Bacterial strains	Erythromycin			Kanamycin			Norfloxacin			Streptomycin			Tetracycline		
	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5
	AG100Atet	64	16 (4) <sup>S</sup>	32 (2) <sup>S</sup>	-	32 (>8) <sup>S</sup>	256	128	16 (8) <sup>S</sup>	128 (1) <sup>I</sup>	<2	<2	<2	2	<2 (>1) <sup>S</sup>
AG102	32	16 (2) <sup>S</sup>	16 (2) <sup>S</sup>	<2	<2	<2	<2	<2	<2	-	128 (>2) <sup>S</sup>	256 (>1) <sup>S</sup>	<2	<2	<2
CM64	-	128 (>2) <sup>S</sup>	256 (>1) <sup>S</sup>	4	<2	<2	4	<2 (>2) <sup>S</sup>	4 (1) <sup>I</sup>	32	4 (8) <sup>S</sup>	8 (4) <sup>S</sup>	nt	nt	nt
KP63	16	<1 (>16) <sup>S</sup>	4 (4) <sup>S</sup>	32	16 (2) <sup>S</sup>	32 (1) <sup>I</sup>	-	128 (>2) <sup>S</sup>	256 (>1) <sup>S</sup>	<4	<4	<4	<2	<2	<2
PA124	128	64 (2) <sup>S</sup>	64 (2) <sup>S</sup>	128	16 (8) <sup>S</sup>	64 (2) <sup>S</sup>	64	8 (8) <sup>S</sup>	32 (2) <sup>S</sup>	nt	nt	nt	8	2 (4) <sup>S</sup>	2 (4) <sup>S</sup>

MIC/2.5: concentration of plant extract added equal to 204.8 µg/mL for AG100A<sub>TET</sub>, KP63; and to 409.6 µg/mL for PA124, CM64, AG102.  
 MIC/5: concentration of plant extract added equal to 102.4 µg/mL for AG100A<sub>TET</sub>, KP63; and to 204.8 µg/mL for PA124, CM64, AG102.  
 (I): Folds decreasing of MIC. S: synergy. I: indifference. nt: not tested; (-): MIC > 256.

18 h [5]. The final concentration of DMSO was lower than 2.5 % and this concentration also served as negative control [5]. Chloramphenicol was used as reference antibiotic. The MICs of samples were detected after 18 h incubation at 37 °C, following addition (40 µl) of 0.2 mg/ml INT and incubation at 37 °C for 30 minutes [5]. MIC was defined as the lowest sample concentration that prevented the color change of the medium and exhibited complete inhibition of microbial growth [27].

Samples were tested alone and then, in the presence of PAβN at 20 mg/L final concentration as previously

reported [5]. Four of the best extracts, those from *A. citratum*, *C. zeylanicum*, *D. psilurus* and *T. tetraptera* were also tested in association [5] at the concentrations selected following a preliminary assay on *P. aeruginosa* PA124 (See Additional file 1: Table S1). All assays were performed in triplicate and repeated thrice. Fractional inhibitory concentration (FIC) [5] were calculated and the interpretations were made as follows: synergistic (<0.5), indifferent (0.5 to 4), or antagonistic (>4) [28] (The FIC values available in Additional file 1: Table S2 and S3).

**Table 5 Minimal inhibitory concentration (MIC) of antibiotics in absence and presence of *Cinnamomum zeylanicum* extract (µg/ml)**

Bacterial strains	Antibiotics and MIC in absence and presence of <i>Cinnamomum zeylanicum</i> extract														
	Ampicillin			Cefepime			Chloramphenicol			Ciprofloxacin			Cloxacillin		
	Alone	MIC/2.5	MIC/5	Alone	MIC/2	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5
AG100Atet	-	-	-	-	-	-	256	16 (16) <sup>S</sup>	32 (8) <sup>S</sup>	256	64 (4) <sup>S</sup>	128 (2) <sup>S</sup>	-	-	-
AG102	-	-	-	128	256 (0.5) <sup>I</sup>	256 (1) <sup>I</sup>	16	8 (2) <sup>S</sup>	16 (1) <sup>I</sup>	<2	<2	<2	-	256 (>1) <sup>S</sup>	-
CM64	-	256 (>1) <sup>S</sup>	-	-	256 (>1) <sup>S</sup>	-	nt	nt	nt	nt	nt	nt	-	-	-
KP63	-	32 (>8) <sup>S</sup>	-	256	32 (8) <sup>S</sup>	256 (1) <sup>I</sup>	-	32 (>8) <sup>S</sup>	256 (>1) <sup>S</sup>	64	128 (0.5) <sup>I</sup>	128(0.5) <sup>I</sup>	-	64 (>4) <sup>S</sup>	256 (>1) <sup>S</sup>
PA124	128	16 (8) <sup>S</sup>	64 (2) <sup>S</sup>	128	128 (1) <sup>I</sup>	128 (1) <sup>I</sup>	32	2 (16) <sup>S</sup>	8 (4) <sup>S</sup>	16	8 (2) <sup>S</sup>	16 (1) <sup>I</sup>	-	-	-
Bacterial strains	Erythromycin			Kanamycin			Norfloxacin			Streptomycin			Tetracycline		
	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5
	AG100Atet	64	16 (4) <sup>S</sup>	32 (2) <sup>S</sup>	-	16 (>16) <sup>S</sup>	128 (>2) <sup>S</sup>	128	128 (1) <sup>I</sup>	128 (1) <sup>I</sup>	<2	<2	<2	2	2 (1) <sup>I</sup>
AG102	32	16 (2) <sup>S</sup>	16 (2) <sup>S</sup>	<2	<2	<2	<2	<2	<2	-	256 (>1) <sup>S</sup>	256 (>1) <sup>S</sup>	<2	<2	<2
CM64	-	128 (>2) <sup>S</sup>	256 (>1) <sup>S</sup>	4	<2 (>2) <sup>S</sup>	<2 (>2) <sup>S</sup>	4	<2 (>2) <sup>S</sup>	4 (1) <sup>I</sup>	32	4 (8) <sup>S</sup>	8 (4) <sup>S</sup>	nt	nt	nt
KP63	16	1 (16) <sup>S</sup>	4 (4) <sup>S</sup>	32	32 (1) <sup>I</sup>	32 (1) <sup>I</sup>	-	128 (>2) <sup>S</sup>	256 (>1) <sup>S</sup>	<4	<4	<4	<2	<2	<2
PA124	128	16 (8) <sup>S</sup>	32 (4) <sup>S</sup>	128	8 (16) <sup>S</sup>	32 (4) <sup>S</sup>	64	32 (2) <sup>S</sup>	64 (1) <sup>I</sup>	nt	nt	nt	8	2 (4) <sup>S</sup>	2 (4) <sup>S</sup>

MIC/2.5: concentration of plant extract added equal to 204.8 µg/mL for AG100A<sub>TET</sub>, KP63; and to 409.6 µg/ml for PA124, CM64, AG102.  
 MIC/5: concentration of plant extract added equal to 102.4 µg/mL for AG100A<sub>TET</sub>, KP63; and to 204.8 µg/ml for PA124, CM64, AG102.  
 (I): Folds decreasing of MIC. S: synergy. I: indifference. nt: not tested; (-): MIC > 256 µg/ml.

**Table 6 Minimal inhibitory concentration (MIC) of antibiotics in absence and presence extracts *Dorstenia psilurus* (µg/ml)**

Bacterial strains	Antibiotics and MIC in absence and presence of <i>Dorstenia psilurus</i> extract														
	Ampicillin		Cefepime		Chloramphenicol		Ciprofloxacin		Cloxacillin						
	Alone	MIC/2.5	MIC/5	Alone	MIC/2	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5			
AG100Atet	-	-	-	-	-	-	256	128 (2) <sup>S</sup>	256 (1) <sup>I</sup>	256	64 (4) <sup>S</sup>	128 (2) <sup>S</sup>	-	-	-
AG102	-	-	-	128	256 (0.5) <sup>I</sup>	256 (0.5) <sup>I</sup>	16	4 (4) <sup>S</sup>	4 (4) <sup>S</sup>	<2	<2	<2	-	-	-
CM64	-	256 (>1) <sup>S</sup>	-	-	64 (>4) <sup>S</sup>	64 (>4) <sup>S</sup>	nt	nt	nt	nt	nt	nt	-	256 (>1) <sup>S</sup>	-
KP63	-	32 (>8) <sup>S</sup>	-	256	64 (4) <sup>S</sup>	128 (2) <sup>S</sup>	-	64 (>4) <sup>S</sup>	256 (>1) <sup>S</sup>	64	64 (1) <sup>I</sup>	64 (1) <sup>I</sup>	-	64 (>4) <sup>S</sup>	256 (>1) <sup>S</sup>
PA124	128	64 (2) <sup>S</sup>	64 (2) <sup>S</sup>	128	128 (1) <sup>I</sup>	128 (1) <sup>I</sup>	32	16 (2) <sup>S</sup>	32 (1) <sup>I</sup>	16	16 (1) <sup>I</sup>	16 (1) <sup>I</sup>	-	-	-

  

Bacterial strains	Antibiotics and MIC in absence and presence of <i>Dorstenia psilurus</i> extract														
	Erythromycin		Kanamycin		Norfloxacin		Streptomycin		Tetracycline						
	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5			
AG100Atet	64	32 (2) <sup>S</sup>	32 (2) <sup>S</sup>	-	128 (>2) <sup>S</sup>	256 (>1) <sup>S</sup>	128	128 (1) <sup>I</sup>	256 (0.5) <sup>I</sup>	<2	<2	<2	2	2 (1) <sup>I</sup>	2 (1) <sup>I</sup>
AG102	32	64 (0.5) <sup>I</sup>	64 (0.5) <sup>I</sup>	<2	<2	<2	<2	<2	<2	-	256 (>1) <sup>S</sup>	256 (>1) <sup>S</sup>	<2	<2	<2
CM64	-	64 (>4) <sup>S</sup>	256 (>1) <sup>S</sup>	4	4 (1) <sup>I</sup>	8 (0.5) <sup>I</sup>	4	<2 (>2) <sup>S</sup>	<2 (>2) <sup>S</sup>	32	8 (4) <sup>S</sup>	32 (1) <sup>I</sup>	nt	nt	nt
KP63	16	1 (16) <sup>S</sup>	8 (2) <sup>S</sup>	32	32 (1) <sup>I</sup>	64 (0.5) <sup>I</sup>	-	-	-	<4	<4	<4	<2	<2	<2
PA124	128	64 (2) <sup>S</sup>	128 (1) <sup>I</sup>	128	4 (32) <sup>S</sup>	16 (8) <sup>S</sup>	64	32 (2) <sup>S</sup>	64 (1) <sup>I</sup>	nt	nt	nt	8	2 (4) <sup>S</sup>	8 (1) <sup>I</sup>

MIC/2.5: concentration of plant extract added equal to 204.8 µg/mL for AG100A<sub>TET</sub>, CM64, KP63, AG102 and to 409.6 µg/ml for PA124.  
 MIC/5: concentration of plant extract added equal to 102.4 µg/mL for AG100A<sub>TET</sub>, CM64, KP63, AG102; and to 204.8 µg/ml for PA124.  
 (): Folds decreasing of MIC. S: synergy. I: indifference. nt: not tested; (-): MIC > 256 µg/ml.

## Results

### Phytochemical composition of the spice extracts

The results of qualitative analysis showed that each plant contains various phytochemicals compounds such as alkaloids, anthocyanins, anthraquinones, flavonoids, phenols, saponins, steroids, tannins and triterpenes as shown in Table 2.

### Antibacterial activity of the spice extracts

The results summarized in Table 3 summarize the MIC of the extract tested alone or in combination with PAβN on the tested microorganisms. It shows that all the studied extracts were active on at least one microbial strain. *A. citratum* showed the best activity, its inhibitory effect being recorded on 85% (24/28) of the tested bacteria. Other

**Table 7 Minimal inhibitory concentration (MIC) of antibiotics in absence and presence extracts *Tetrapleura tetraptera* (µg/ml)**

Bacterial strains	Antibiotics and MIC in absence and presence <i>Tetrapleura tetraptera</i> extract														
	Ampicillin		Cefepime		Chloramphenicol		Ciprofloxacin		Cloxacillin						
	Alone	MIC/2.5	MIC/5	Alone	MIC/2	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5			
AG100Atet	-	-	-	-	-	-	256	256 (1) <sup>I</sup>	-	256	128 (2) <sup>S</sup>	128 (2) <sup>S</sup>	-	-	-
AG102	-	-	-	128	256 (0.5) <sup>I</sup>	256 (0.5) <sup>I</sup>	16	8 (2) <sup>S</sup>	8 (2) <sup>S</sup>	<2	<2	<2	-	-	-
CM64	-	-	-	-	-	-	nt	nt	nt	nt	nt	nt	-	-	-
KP63	-	-	-	256	256 (1) <sup>I</sup>	-	-	256 (>1) <sup>S</sup>	256 (>1) <sup>S</sup>	64	64 (1) <sup>I</sup>	64 (1) <sup>I</sup>	-	128 (>2) <sup>S</sup>	256 (>1) <sup>S</sup>
PA124	128	64 (2) <sup>S</sup>	128 (1) <sup>I</sup>	128	128 (1) <sup>I</sup>	128 (1) <sup>I</sup>	32	4 (8) <sup>S</sup>	8 (4) <sup>S</sup>	16	16 (1) <sup>I</sup>	16 (1) <sup>I</sup>	-	-	-

  

Bacterial strains	Antibiotics and MIC in absence and presence of <i>Tetrapleura tetraptera</i> extract														
	Erythromycin		Kanamycin		Norfloxacin		Streptomycin		Tetracycline						
	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5	Alone	MIC/2.5	MIC/5			
AG100Atet	64	64 (1) <sup>I</sup>	64 (1) <sup>I</sup>	-	256 (>1) <sup>S</sup>	256 (>1) <sup>S</sup>	128	128 (1) <sup>I</sup>	256 (0.5) <sup>I</sup>	<2	<2	<2	2	2 (1) <sup>I</sup>	2 (1) <sup>I</sup>
AG102	32	64 (0.5) <sup>I</sup>	64 (0.5) <sup>I</sup>	<2	<2	<2	<2	<2	<2	-	256 (>1) <sup>S</sup>	256 (>1) <sup>S</sup>	<2	<2	<2
CM64	-	256 (>1) <sup>S</sup>	-	4	4 (1) <sup>I</sup>	8 (0.5) <sup>I</sup>	4	4 (1) <sup>I</sup>	8 (0.5) <sup>I</sup>	32	16 (2) <sup>S</sup>	32 (1) <sup>I</sup>	nt	nt	nt
KP63	16	<1 (>16) <sup>S</sup>	8 (2) <sup>S</sup>	32	32 (1) <sup>I</sup>	64 (0.5) <sup>I</sup>	-	256 (>1) <sup>S</sup>	-	<4	<4	<4	<2	<2	<2
PA124	128	64 (2) <sup>S</sup>	64 (2) <sup>S</sup>	128	64 (2) <sup>S</sup>	64 (2) <sup>S</sup>	64	32 (2) <sup>S</sup>	64 (1) <sup>I</sup>	nt	nt	nt	8	2 (4) <sup>S</sup>	2 (4) <sup>S</sup>

MIC/2.5: concentration of plant extract added equal to 204.8 µg/mL for CM64 and to 409.6 µg/ml for AG100A<sub>TET</sub>, PA124, KP63, AG102.  
 MIC/5: concentration of plant extract added equal to 102.4 µg/mL for CM64; and to 204.8 µg/ml for AG100A<sub>TET</sub>, PA124, KP63, AG102.  
 (): Folds decreasing of MIC. S: synergy. I: indifference. nt: not tested; (-): MIC > 256 µg/ml.

samples were less active, their inhibitory potencies being observed on 75% of tested bacteria (21/28) for *I. cylindricum* and *C. zeylanicum*, 67.9 % (19/28) for *A. melegueta*, *D. psilurus*, *F. lepieuri* and *T. tetraptera*; 64.3% (18/28) for *M. myristica* and *S. zenkeri*; 50 % (14/28) for *P. guineense* and 42.9 % (12/28) for *P. brazzeana*.

#### **Role of efflux pumps in susceptibility of gram negative bacteria to the tested spice extracts**

Potentiating effect of EPI was not observed on tested bacteria when associated with *M. myristica*, *P. brazzeana*, *T. tetraptera* and *S. zenkeri*. PA $\beta$ N weakly increased the activity of *A. citratum*, *A. melegueta*, *F. lepieuri*, *I. cylindricum*, *C. zeylanicum* and *P. guineense*. The activity of *D. psilurus* in the presence of EPI significantly increased on most of the tested bacteria (except against *P. stuartii* ATCC29916, *E. cloacae* ECC169 and *E. aerogenes* EA27) (see Table 3).

#### **Effects of the association of some spice extracts with antibiotics**

*A. citratum*, *C. zeylanicum*, *D. psilurus* and *T. tetraptera* (Tables 4, 5, 6 and 7) were associated to antibiotics in view of evaluating the possible synergistic effect of these associations. A preliminary study using *P. aeruginosa* PA124 was carried out with ten antibiotics (CLX, AMP, ERY, KAN, CHL, TET, FEP, STR, CIP and NOR) to select the appropriate sub-inhibitory concentrations to be used. MIC/2.5 and MIC/5 were then selected as the sub-inhibitory concentrations (see Additional file 1: Table S1). All of these four extracts were then tested in association with antibiotics previously listed on strains of *E. coli* AG100A<sub>TET</sub> and AG102, *E. aerogenes* CM64, *K. pneumoniae* KP63 and *P. aeruginosa* PA124. No antagonistic effect (FIC > 4) was observed between extracts and antibiotics meanwhile indifference was observed between *T. tetraptera* and antibiotics in most of the case (see Tables 5, 6, and 7, Additional file 1: S2, S3, S4 and S5). Significant increase of the activity was observed with the association of the extracts of *A. citratum* and *D. psilurus* on *E. aerogenes* CM64 and *K. pneumoniae* KP63, and with *C. zeylanicum* against *K. pneumoniae* KP63. A significant decrease (synergy effect) of MIC values was also observed when ERY was associated with various extracts, and when extracts of *A. citratum* and *C. zeylanicum* were each combined with aminoglycosides (KAN, STR), the best activity being noted against *E. aerogenes* CM64.

## **Discussion**

### **Phytochemical composition of the spice extracts**

The phytochemical studies revealed the presence of secondary metabolite such as alkaloids, anthocyanins, anthraquinones, flavonoids, phenols, saponins, sterols, tannins and triterpenes; several molecules belonging to these classes of secondary metabolites were found active on pathogenic microorganisms [29].

### **Antibacterial activity of the spice extract**

Although this is the first time that plants used in this work are studied for their activities vis-à-vis multi-resistant bacteria, plants belonging to some of the genus studied herein, like the *Aframomum* genus are well documented for their antimicrobial activity [6]. Some antibacterial compounds, such as acridone and chelerythrine have previously been isolated from the fruits of *F. lepieurii* [14,30]. The antimicrobial activity of *P. brazzeana* and *S. zenkeri* is mainly due to some sulfur compounds. In fact, sulfur compounds with antimicrobial properties have previously been isolated from the two plants [7,31]. Several alkaloids of the genus *Piper* proved to be responsible for the activity of *P. guineense* [32]. The detection of this class of secondary metabolites in the extract studied herein can explain the observed activities. According to Krishnaiah et al. [16], the antimicrobial activity of *I. cylindricum* can be due to the presence of tannins in this plant. However, tannins were not detected in the extract of *I. cylindricum* as found in the present work (Table 2), suggesting that other classes of secondary metabolites might be responsible for the antibacterial activity of this plant.

### **Role of efflux pumps in susceptibility of gram negative bacteria to the tested spice extracts**

The significant increase of the activity of the extract of *D. psilurus* in the presence of EPI, indicates that bioactive constituents of this plant extract are substrate of efflux pumps. Efflux through AcrAB-TolC pumps was reported as essential mode of resistance of several Gram-negative MDR bacteria to a number of flavonoids isolated from plants of the genus *Dorstenia*, such as isobavachalcone, kanzonol C, stipulin, etc. [4,15,33-35]. This suggests that possible combination of the extract of *D. psilurus* with EPI can be envisaged to overcome MDR bacteria.

### **Effects of the association of extracts with antibiotics**

The results obtained by combining the antibiotic with the extracts of *A. citratum*, *C. zeylanicum*, *D. psilurus* and *T. tetraptera* indicate that these extracts contain chemical compounds that can modulate the activity of antibiotics against bacteria expressing MDR phenotypes. The methanol extracts of *A. citratum*, *C. zeylanicum* and *D. psilurus* showed a synergistic effect with antibiotics inhibiting bacterial cell wall synthesis (AMP and CEF) on *K. pneumoniae* KP63. The intrinsic mode of action of the active extracts is to be investigated.

## **Conclusion**

The present work evidently provides information in the role of some Cameroonian spices in the fight against multi-resistant bacteria. The study also highlights the potential of *D. psilurus* as a strong antibacterial agent

when the extract is combined with efflux pump inhibitor and several antibiotics.

## Additional file

**Additional file 1: Table S1.** Activities of antibiotics in combination with the sub-inhibitory concentrations of some plants extracts on *Pseudomonas aeruginosa* PA124. **S2.** Fractional inhibitory Concentrations of the association between antibiotics and extracts of *Aframomum citratum* at MIC/2.5 and MIC/5 (µg/ml) against MDR bacteria. **S3.** Fractional inhibitory Concentrations of the association between antibiotics and extracts of *Cinnamomum zeylanicum* at MIC/2.5 and MIC/5 (µg/ml) against MDR bacteria. **S4.** Fractional inhibitory Concentrations of the association between antibiotics and extracts of *Dorstenia ptilurus* at MIC/2.5 and MIC/5 (µg/ml) against MDR bacteria. **S5.** Fractional inhibitory Concentrations of the association between antibiotics and extracts of *Tetrapleura tetraptera* at MIC/2.5 and MIC/5 (µg/ml) against MDR bacteria.

## Competing interest

The authors declare that they have no competing interest.

## Authors' contributions

IKV carried out the study; VK designed the experiments and wrote the manuscript; VK, GAF, JAKN, JPD, JRK and JMP supervised the work; VK and JMP provided the bacterial strains; All authors read and approved the final manuscript.

## Acknowledgements

Authors are thankful to the Cameroon National Herbarium (Yaounde) for plants identification, Mr Simplicie R. Mouokeu for its technical support, and Mr. Paul K. Lunga for language editing.

## Author details

<sup>1</sup>Department of Biochemistry, Faculty of science, University of Dschang, P.O. Box 67, Dschang, Cameroon, Africa. <sup>2</sup>Transporteurs Membranaires, Chimiorésistance et Drug Design, UMR-MD1, IFR 88, UFRs de Médecine et de Pharmacie, Marseille, France.

Received: 27 January 2012 Accepted: 4 June 2012

Published: 15 June 2012

## References

- Ziegler J: *L'Empire de la honte*. Paris, France: Ed Fayard; 2005.
- OCDE: *Partenariat contre les maladies infectieuses*. 2009 2009. [http://www.observeurocde.org/news/fullstory.php/aid/743/Partenariat\\_contre\\_les\\_maladies\\_infectieuses.html](http://www.observeurocde.org/news/fullstory.php/aid/743/Partenariat_contre_les_maladies_infectieuses.html), 2009. Accessed on 22 February 2010.
- Yala D, Merad AS, Mohamedi D, Ouar Korich MN: **Classification et mode d'action des antibiotiques**. *Médecine du Maghreb* 2001, **91**:5–12.
- Kuete V, Krusche B, Youns M, Voukeng I, Fankam AG, Tankeo S, Lacmata S, Efferth T: **Cytotoxicity of some Cameroonian spices and selected medicinal plant extracts**. *J Ethnopharmacol* 2011, **134**(3):803–812.
- Fankam AG, Kuete V, Voukeng IK, Kuete JR, Pages JM: **Antibacterial activities of selected Cameroonian spices and their synergistic effects with antibiotics against multidrug-resistant phenotypes**. *BMC Complement Altern Med*. 2011, **11**:104.
- Titanji VPK, Zofou D, Ngemenya MN: **The antimalarial potential of medicinal plants used for the treatment of malaria in Cameroonian folk medicine**. *Afr J Trad CAM* 2008, **5**(3):302–321.
- Ngemenya MN, Mbah JA, Tane P, Titanji VPK: **Antibacterial effects of some Cameroonian medicinal plants against common pathogenic bacteria**. *Afr J Trad CAM* 2006, **3**(2):84–93.
- Konning GH, Agyare C, Ennon B: **Antimicrobial activity of some medicinal plants from Ghana**. *Fitoterapia* 2004, **75**(1):65–67.
- Okigbo RN, Ogbonnaya UO: **Antifungal effects of two tropical plant leaf extracts (*Ocimum gratissimum* and *Aframomum melegueta*) on postharvest yam (*Dioscorea* spp.) rot**. *Afr J Biotechnol* 2006, **5**(9):727–731.
- Ranasinghe L, Jayawardena B, Abeywickrama K: **Fungicidal activity of essential oils of *Cinnamomum zeylanicum* (L.) and *Syzygium aromaticum* (L.) Merret L. M. Perry against crown rot and anthracnose pathogens isolated from banana**. *Lett Appl Microbiol* 2002, **35**(3):208–211.
- Shahverdi AR, Monsef-Esfahani HR, Tavasoli F, Zaheri A, Mirjani R: **Trans-Cinnamaldehyde from *Cinnamomum zeylanicum* bark essential oil reduces the Clindamycin resistance of *Clostridium difficile* in vitro**. *J Food Sci* 2007, **72**(1):55–58.
- Abegaz BM, Ngadjui BT, Dongo E, Bezabiha MT: **Chemistry of the Genus *Dorstenia***. *Curr Org Chem* 2000, **4**(10):1079–1090.
- Dimo T, Rakotonirina A, Tan PV, Dongo E, Dongmo AB, Kamtchoung P, Azay J, Abegaz BM, Cros G, Ngadjui TB: **Antihypertensive effects of *Dorstenia ptilurus* extract in fructose-fed hyperinsulinemic, hypertensive rats**. *Phytomedicine* 2001, **8**(2):101–106.
- Ngono AN, Biyiti L, Amvam Zollo PH, Bouchet P: **Evaluation of antifungal activity of extracts of two Cameroonian Rutaceae: *Zanthoxylum lepreurii* Guill. et Perr., and *Zanthoxylum xanthoxyloides* Watterm.** *J Ethnopharmacol* 2000, **70**(3):335–342.
- Kuete V: **Potential of Cameroonian plants and derived products against microbial infections: a review**. *Planta Med* 2010, **76**(14):1479–1491.
- Krishnaiah D, Devi T, Bono A, Sarbaty R: **Studies on phytochemical constituents of six Malaysian medicinal plants**. *J Med Plant Res* 2009, **3**(2):67–72.
- Nishimoto K, Ito M, Natori S, Ohmoto T: **The structures of arundoin, cylindrin and fernenoltriterpenoids of fernane and arborane of *Imperata cylindrica* var. *koenigif.*** *Tetrahydron* 1968, **24**:735–752.
- Nguefack J, Letha V, Amvam Zollo PH, Mathur SB: **Evaluation of five essential oils from aromatic plants of Cameroon for controlling food spoilage and mycotoxin producing fungi**. *Int J Food Microbiol* 2004, **94**(3):329–334.
- Burubai W, Akor AJ, Igoni AH, Puyate YT: **Some physical properties of African nutmeg (*Monodora myristica*)**. *Int Agrophysics* 2007, **21**:123–126.
- Noumi E, Dibakto TW: **Medicinal plants used for peptic ulcer in the Bangangte region, western Cameroon**. *Fitoterapia* 2000, **71**(4):406–412.
- Oparaeke AM, Bunmi JO: **Bioactivity of two powdered spices *Piper guineense* thonn & schum and *Xylopia aethiopica* (dunal) a. Richard] as home-made insecticides against *Callosobruchus subinnotatus* (pic.) on stored bambara groundnut**. *Agric Trop Subtrop* 2006, **39**(2):132–134.
- Nwinyi OC, Chinedu NS, Ajani OO, Ikpo CO, Ogunniran KO: **Antibacterial effects of extracts of *Ocimum gratissimum* and *Piper guineense* on *Escherichia coli* and *Staphylococcus aureus***. *Afr J Food Sci* 2009, **3**(3):77–81.
- Kouokam JC, Jahns T, Becker H: **Antimicrobial Activity of the Essential Oil and Some isolated Sulfur-Rich Compounds from *Scorodophloeus zenkeri***. *Planta Med* 2002, **68**(12):1082–1087.
- Harborne JB: *Phytochemical Methods*. New York, USA: Chapman and Hall; 1973.
- Eloff JN: **A sensitive and quick microplate method to determine the minimal inhibitory concentration of plant extracts for bacteria**. *Planta Med* 1998, **64**(8):711–713.
- Mativandlela SPN, Lall N, Meyer JJM: **Antibacterial, antifungal and antitubercular activity of (the roots of) *Pelargonium reniforme* (CURT) and *Pelargonium sidoides* (DC) (Geraniaceae) root**. *S Afr J Bot* 2006, **72**(2):232–237.
- Kuete V, Ngameni B, Fotso Simo CC, Kengap Tankeu R, Tchaleu Ngadjui B, Meyer JJM, Lall N, Kuete JR: **Antimicrobial activity of the crude extracts and compounds from *Ficus chlamydocarpa* and *Ficus cordata* (Moraceae)**. *J Ethnopharmacol* 2008, **120**(1):17–24.
- Coutinho HD, Vasconcellos A, Freire-Pessôa HL, Gadelha CA, Gadelha TS, Almeida-Filho GG: **Natural products from the termite *Nasutitermes corniger* lower aminoglycoside minimum inhibitory concentrations**. *Pharmacogn Mag* 2010, **6**(21):1–4.
- Cowan MM: **Plant products as antimicrobial agents**. *Clin Microbiol Rev* 1999, **12**(4):564–582.
- Adesina SK: **The Nigerian *Zanthoxylum*; chemical and biological values**. *Afr J Trad CAM* 2005, **2**(3):282–301.
- El Migirab S, Bergbr Y, Jadot J: **Isothiocyanates, Thiourees et Thiocarbamates isolés de *Pentadiplandra brazzeana***. *Phytochemistry* 1977, **16**(11):1719–1721.
- Parmar VS, Jain SC, Bisht KS, Jain R, Taneja P, Jha A, Tyagi OD, Prasad AK, Wengel J, Olsen CE, Bol PM: **Phytochemistry of the genus *Piper***. *Phytochemistry* 1997, **46**(4):591–673.

33. Omisore NOA, Adewunmi CO, Iwalewa EO, Ngadjui BT, Adenowo TK, Abegaz BM, Ojewole JA, Watchueng J: **Antitrichomonal and antioxidant activities of *Dorstenia barteri* and *Dorstenia convexa***. *Braz J Med Biol Res* 2005, **38**(7):1087–1094.
34. Ngameni B, Watchueng J, Fekam BF, Keumedjio F, Ngadjui TB, Gut J, Abegaz BM, Rosenthal PJ: **Antimalarial prenylated chalcones from the twigs of *Dorstenia barteri* var. *subtriangularis***. *ARKIVOC* 2007, **13**:116–123.
35. Kuete V, Ngameni B, Tangmouo GJ, Bolla JM, Alibert-Franco S, Ngadjui TB, Pagès JM: **Efflux Pumps Are Involved in the Defense of Gram-Negative Bacteria against the Natural Products Isobavachalcone and Diospyrone**. *Antimicrob Agents Chemother* 2010, **54**(5):1749–1752.

doi:10.1186/1756-0500-5-299

**Cite this article as:** Voukeng *et al.*: Antibacterial and antibiotic-potential activities of the methanol extract of some cameroonian spices against Gram-negative multi-drug resistant phenotypes. *BMC Research Notes* 2012 **5**:299.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

