

CORRECTION

Open Access



Correction to: High rhesus (Rh(D)) negative frequency and ethnic-group based ABO blood group distribution in Ethiopia

Lemu Golassa^{1†}, Arega Tsegaye^{2†}, Berhanu Erko¹ and Hassen Mamo^{3*}

Correction to: *BMC Res Notes* (2017) 10:330

<https://doi.org/10.1186/s13104-017-2644-3>

Following publication of the original article [1] the authors reported that the information in Ref. [32] had been misquoted. The Rh factor in one region in Saudi Arabia is 8.8%, not 29% as stated.

Author details

¹ Aklilu Lemma Institute of Pathobiology, Addis Ababa University, P.O. Box 1176, Addis Ababa, Ethiopia. ² Jimma University, P.O. Box 378, Jimma, Ethiopia. ³ Department of Microbial, Cellular and Molecular Biology, College of Natural Sciences, Addis Ababa University, P.O. Box 1176, Addis Ababa, Ethiopia.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 7 March 2018 Accepted: 7 March 2018

Published online: 14 March 2018

Reference

1. Golassa L, Tsegaye A, Erko B, Mamo H. High rhesus (Rh(D)) negative frequency and ethnic-group based ABO blood group distribution in Ethiopia. *BMC Res Notes*. 2017;10:330. <https://doi.org/10.1186/s13104-017-2644-3>.

The original article can be found online at <https://doi.org/10.1186/s13104-017-2644-3>.

*Correspondence: binmamo@yahoo.com; hassen.mamo@aau.edu.et

[†]Lemu Golassa and Arega Tsegaye have contributed equally to this work

³ Department of Microbial, Cellular and Molecular Biology, College of Natural Sciences, Addis Ababa University, P.O. Box 1176, Addis Ababa, Ethiopia

Full list of author information is available at the end of the article